

UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



AGROMET-ADVISORY BULLETIN

Date: 07.10.2022

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department The forecast is valid for Bengaluru Rural district Weather forecast (Valid from 08-10-2022 to 12-10-2022)

Forecast summary:

| Parameters | 08.10.2022 | 09.10.2022 | 10.10.2022 | 11.10.2022 | 12.10.2022 |
|----------------------------------|------------|------------|------------|------------|------------|
| Rainfall (mm) | 4 | 5 | 15 | 15 | 5 |
| Max Temp Trend (^o C) | 27 | 26 | 27 | 27 | 27 |
| Min Temp Trend (°C) | 20 | 20 | 20 | 20 | 20 |
| Total cloud cover (octa) | 7 | 7 | 8 | 8 | 7 |
| Relative humidity (%)Max | 86 | 86 | 88 | 88 | 87 |
| Relative humidity (%)Min | 74 | 74 | 76 | 76 | 75 |
| Wind speed(Km/hr) | 6 | 7 | 6 | 7 | 6 |
| Wind Direction (Degrees) | 290 | 292 | 248 | 212 | 212 |

Light to moderate rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 26.0-27.0°C and Minimum of 20.0°C. Relative humidity 86-88 % during morning hrs and 74-76 % during noon is expected. Wind speed is 6-7 km/hr.

Weather Based Agro Advisories

Crop information and Crop Stages of the major Kharif/Rabi crops

| District | Kharif crops Hort | | | Horticul | ture crops | |
|------------------------|-------------------|---------|----------------------|--------------|------------|-------|
| Bangalore Rural | Groundnut | Redgram | Finger millet | Maize | Grape | Mango |
| (BR) | PF,PM | V,F | V,T,E | Ts, CI,GF,FD | - | |

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching, CI: Cob Initiation

Agromet Advisory:

| Crop/ Component | Stage/ Condition | Pest and Disease | Agro advisories |
|--------------------|---------------------|--|-----------------|
| General | | 1. Hasta rainstar starts from September 27 th and remains up to October 10 th . The normal rainfall of Hasta rainstar is 105.8 mm. | |

| Kharif | Vegetative | The following crops are suggested for sowing. | | |
|--------|---|--|--|--|
| crops | stage And Tillering stage | Finger millet Ragi (Indaf-5, 7,9, ML-365, KMR-301), Horse gram-PHG-9, KBH-1, Niger, Field bean-HA-3 and 4 and sunflower-KBSH-1,41,42, 44 & 53, cowpea (KBC-1, TVX-944 and PKB-4 for vegetable purpose). 1. Suggested for intercultivation for Ragi crop 2. Wherever over population of crop plants is existing, undertake thinning operation for better availability of scarce soil moisture to the reduced population of crop plants. 3. To control of leaf folder insect in Redgram to spray Methomyl 40SP , 2 gram/litre of water before flowering of crop. 4. If Wilt diseases noticed in Redgram field –Drenching Carbendazim 50 WP 2 g/litre of water. | | |
| | | 5. Remove and burn the infected Wilt diseases Redgram plant in field. Horticulture crop | | |
| Mango | | Paclobutrazol soil application @ 5 g a.i per plant is recommended for uniform flowering behavior in Mango crop. Paclobutrazol is effective not only in flower induction but also in early and off season flower induction which thereby maintains regularity and synchronization in flowering. Sericulture | | |
| | 1. Maintai | in the optimum room temperature in Sericulture unit. | | |
| Dairy | Congen Piggery | Congenial climate and abundant fodder availability facilitate improved Dairy, Piggery, Poultry, Sheep and Goat rearing. | | |
| | Feeding of lactating cow | | | |
| | maximu A concession sourcess as brand Mineral level of | feeding of dairy cattle should envisage minimum wastage of nutrients and um returns in respect of milk produced. entrate mixture made up of protein supplements such as oil cakes, energy such as cereal grains (Maize, Jowar), tapioca chips and laxative feeds such s (rice bran, wheat bran, gram husk) is generally used. I mixture containing major and all the trace elements should be included at a 2 percent. fiber for milking cow should be 17 % and NDF (Neutral Detergent Fiber) | | |

AMFU of IMD Bengaluru

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*