

# UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



# AGROMET-ADVISORY BULLETIN

Date: 11.10.2022

#### Issued jointly by, UAS, Bengaluru & Indian Meteorological Department The forecast is valid for Bengaluru Rural district Weather forecast (Valid from 12-10-2022 to 16-10-2022)

Forecast summary:

Parameters	12.10.2022	13.10.2022	14.10.2022	15.10.2022	16.10.2022
Rainfall (mm)	10	17	10	9	9
Max Temp Trend ( <sup>o</sup> C)	27	27	27	27	27
Min Temp Trend ( <sup>o</sup> C)	20	20	20	20	20
Total cloud cover (octa)	8	8	8	8	8
Relative humidity (%)Max	91	91	92	92	92
Relative humidity (%)Min	70	70	71	70	72
Wind speed(Km/hr)	6	7	8	7	6
Wind Direction (Degrees)	202	334	207	68	112

Light to moderate rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 27.0°C and Minimum of 20.0°C. Relative humidity 91-92 % during morning hrs and 70-72 % during noon is expected. Wind speed is 6-7 km/hr.

# Weather Based Agro Advisories

#### Crop information and Crop Stages of the major Kharif/Rabi crops

District	Kharif crops			Horticulture crops		
<b>Bangalore Rural</b>	Groundnut	Redgram	<b>Finger millet</b>	Maize	Grape	Mango
( <b>BR</b> )	PF,PM	V,F	V,T,E	Ts, CI,GF,FD	-	

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching, CI: Cob Initiation

#### Agromet Advisory:

Crop/	Stage/ Condition	Pest and	Agro advisories		
Component		Disease			
General		1. Chitta rainstar starts from October 11 <sup>th</sup> and remains up to			
		October 23 <sup>rd</sup> . The normal rainfall of Chitta rainstar is 55.4			
		mm.			
Kharif	Tillering stage/Ear	The following crops are suggested for sowing.			
crops	head formation Finger millet Ragi (Indaf-5, 7,9, ML-365, KMR-301), Hors				
	stage/flowering/pod	gram- PHG-9, KBH-1, Niger, Field bean-HA-3 and 4 and			
	formation and	sunflower-KBSH-1,41,42, 44 & 53, cowpea (KBC-1, TVX-944 and			
	maturity stage	PKB-4 for vegetat			
		1. Suggested for intercultivation for Ragi crop			
		2. To contro	l of leaf folder insect in Redgram to spray		
		Methomyl	40SP , 2 gram/litre of water before flowering of		
		crop.			
		3. If Wilt d	iseases noticed in Redgram field –Drenching		
			im 50 WP 2 g/litre of water.		
		4. Remove ar	d burn the infected Wilt diseases Redgram plant in		

	field.				
Horticulture crop					
Mango	1. Paclobutrazol soil application @ 5 g a.i per plant is recommended for uniform flowering behavior in Mango crop. Paclobutrazol is effective not only in flower induction but also in early and off season flower induction which thereby maintains regularity and synchronization in flowering.				
	Sericulture				
	1. Maintain the optimum room temperature in Sericulture unit.				
Dairy	Congenial climate and abundant fodder availability facilitate improved Dairy, Piggery, Poultry, Sheep and Goat rearing.				
	Feeding of lactating cow				
	<ul> <li>Proper feeding of dairy cattle should envisage minimum wastage of nutrients and maximum returns in respect of milk produced.</li> <li>A concentrate mixture made up of protein supplements such as oil cakes, energy</li> </ul>				
	sources such as cereal grains (Maize, Jowar), tapioca chips and laxative feeds such as brans (rice bran, wheat bran, gram husk) is generally used.				
	Mineral mixture containing major and all the trace elements should be included at a level of 2 percent.				
	1. Dietary fiber for milking cow should be 17 % and NDF (Neutral Detergent Fiber ) 22 %				

AMFU of IMD Bengaluru

**Important Note:** Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*