



UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU  
GRAMIN KRISHI MAUSAM SEWA(GKMS)  
AMFU OF IMD, BENGALURU



**AGROMET-ADVISORY BULLETIN**

Date: **13.12.2022**

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

**The forecast is valid for Bengaluru Rural district**  
**Weather forecast (Valid from 14-12-2022 to 18-12-2022)**

Forecast summary:

Parameters	14.12.2022	15.12.2022	16.12.2022	17.12.2022	18.12.2022
Rainfall (mm)	16	0	0	0	0
Max Temp Trend ( °C)	27	27	29	28	28
Min Temp Trend ( °C)	17	17	17	17	16
Total cloud cover (octa)	3	7	7	5	6
Relative humidity (%)Max	95	97	77	76	89
Relative humidity (%)Min	69	66	40	43	48
Wind speed(Km/hr)	19	13	10	12	16
Wind Direction (Degrees)	112	112	90	68	68

**Light to medium rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 27.0-29.0°C and Minimum of 16.0-17.0°C. Relative humidity 77-97 % during morning hrs and 40-69 % during noon is expected. Wind speed is 10-19 km/hr.**

**Weather Based Agro Advisories**

**Crop information and Crop Stages of the major Kharif/Rabi crops**

District	Kharif crops				Horticulture crops	
Bangalore Rural (BR)	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
	--	GF,M,H	H	--	-	--

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Transplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting, IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching, CI: Cob Initiation

**Agromet Advisory:**

Crop/Component	Stage/Condition	Pest and Disease	Agro advisories
General			<ul style="list-style-type: none"> <li>Moola rainstar starts from December 16<sup>th</sup> to December 28<sup>th</sup>. The normal rainfall of Moola rainstar is 4.1 mm. <b>The following crops are suggested for sowing.</b></li> <li>Field bean-HA-3 and 4 and sunflower-KBSH-1,41,42, 44 &amp; 53, cowpea (KBC-1, TVX-944 and PKB-4 for vegetable purpose). Horse gram- PHG-9, KBH-1</li> <li>The grains of the harvested crops should be properly dried by retaining moisture percentage of Cereals 11-12 %, Pulses-9%, Oilseeds-8% and Vegetable seeds 5-6% for long storage &amp; also minimize the store pest damage.</li> <li>To protect the pulse grains from storage pests apply oils of Castor/ linseed/honge/neem oil @ 3-5 ml per kg of grains.</li> </ul>
Finger millet	Harvesting		1. Advised for postpone harvesting next couple of days due to cloudy and light rainfall forecasted from IMD

<b>Maize</b>	Harvesting	1. Advised for postpone harvesting next couple of days due to cloudy and light rainfall forecasted from IMD
<b>Horticulture crop</b>		
<b>Mango</b>	Flower bud initiation and Flowering	<ol style="list-style-type: none"> <li>1. Sudden drop in minimum temperature is observed in Mango it will be affects the floral induction and spray pacloburtrazol as plant growth retardant which restrict the vegetative growth.</li> <li>2. Clear the weeds in Mango orchard/Guava/Sapota put it under the basin as mulch.</li> <li>3. If Phanerogamic plants are growing on the mango tree to cut/prune out completely and apply Bordeaux paste or Copper oxy chloride that portion.</li> <li>4. Remove the weeds such as lantana which are growing under the mango tree.</li> <li>5. Leaf hopper and Powdery mildew disease incidence is more before flowering and immediately after fruit formation to manage spraying of Carbaryl, 50WP @4g/litre of water or Imidachlorprid @ 0.3ml/ litre of water for management of leaf hopper.</li> <li>6. Spray Lamda Cyhalothrin 5EC @ 0.5 ml/ litre of water or sulphur dust (SULTAF) 80 W @3g/litre of water against the Powdery mildew diseases.</li> <li>7. If the incidence of Leaf hopper is severe spray Azadirachtin (10,000 ppm) @ 7.0 ml/ litre of water.</li> </ol>
<b>Animal Husbandry</b>		
		<ol style="list-style-type: none"> <li>1. To protect animals from a sudden drop in temperature, keep the animals in a covered shed/area during the night. The bedding/hay in the animal sheds must be kept dry and changed/aired every day.</li> <li>2. Due care should be taken to store/procure fodder for periods of shortage that may occur during the winter months in certain areas. Perennial grasses must be cut at this time.</li> </ol>
<b>Sericulture</b>		
		<ol style="list-style-type: none"> <li>1. White muscardine: caused by <i>Beauveria bassiana</i> ,</li> <li>2. Manage the humidity in the rearing house by providing good cross-ventilation. Dust dry slaked lime powder when silkworms settle for moult.</li> <li>3. Feed silkworms with adequate quantity of mulberry leaves to avoid the accumulation of left over leaves in the rearing bed. Make sure that the silkworm bed is dry and thin.</li> <li>4. If the silkworm rearing house temperature falls below 22°C, raise it using room heater / charcoal stove.</li> <li>5. Collect muscardine affected larvae from the rearing bed before mummification, dust antimuscardine bed disinfectant and finally burn them. Do not throw them on the street or feed to animals / birds.</li> <li>6. Dust Vijetha and Vijetha Supplement or Ankush bed disinfectant as per recommended schedule or dust any recommended anti-muscardine bed disinfectant as per the schedule.</li> </ol>
<b>Poultry</b>		
		<ol style="list-style-type: none"> <li>1. The poultry house should be located in such a way that long axis is in east-west direction. This will prevent the direct sunshine over the birds.</li> <li>2. Beginning at one day of age, the chick should be housed at a temperature 35° C will maintain one week, at a relative humidity between 40 – 60% after wards 2 to 4 weeks temperature decreases every week by 2° C.</li> </ol>

- |  |   |
|--|---|
|  | <ol style="list-style-type: none"><li>3. Provide artificial brooding to chicks to maintain adequate temperature.</li><li>4. Care should be taken to prevent the chicks from being exposed to wind chill.</li><li>5. Sides should be covered with curtains during cool hours of the day.</li><li>6. Wet litter material should be removed regularly</li><li>7. Ensure proper cross ventilation to avoid ammonia accumulation</li></ol> |
|--|---|

**AMFU of IMD  
Bengaluru**

**Important Note:** Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website:  
*mausam.imd.gov.in*