

UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



AGROMET-ADVISORY BULLETIN

Date: 04.07.2023

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

The forecast is valid for Bengaluru urban district.

Significant past weather for the preceding week					
Parameter	30.06.2023	01.07.2023	02.07.2023	03.07.2023	04.07.2023
Rainfall (mm)	3.2	0	0	0.8	2.2
Max. temp(°C)	30.0	29.8	29.8	28.2	30.2
Min.Temp(°C)	19.8	20.6	21.4	19.0	20.0
Sky condition(Octas)	4	2	4	6	4
Relative humidity(%) 0830 hours	83	88	84	83	76
Relative humidity(%) 1730 hours	58	51	54	44	
Wind Speed (kmph)	6.5	8.0	6.6	5.0	8.4
Wind Direction	230	230	320	230	230

Weather forecast (Valid from 05-07-2023 to 09-06-2023)

Forecast summary:

Parameters	05.07.2023	06.07.2023	07.07.2023	08.07.2023	09.07.2023
Rainfall (mm)	14	13	11	4	3
Max Temp Trend (^o C)	30	29	28	28	28
Min Temp Trend (^o C)	21	20	20	20	20
Total cloud cover (octa)	6	6	6	6	4
Relative humidity (%)Max	89	89	87	87	87
Relative humidity (%)Min	85	85	84	84	84
Wind speed(Km/hr)	10	9	11	12	10
Wind Direction (Degrees)	248	248	248	248	248

Light to medium rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 28.0-30.0°C and Minimum of 20.0-21.0°C. Relative humidity 87-89 % during morning hrs and 84-85 % during noon is expected. Wind speed is 9-12 km/hr. Weather Based Agro Advisories

Cron information and Cron Stages of the major Kharif/Rabi crons

Crop mormation and Crop Stages of the major Kharn/Kabi crops						
District	Kharif crops Horticulture crops				ture crops	
Bangalore	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
Urban (BU)	S.G,EV	S,G,EV	-	S,G,EV	-	-

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

Agromet Advisory:

Crop/	Stage/	Pest and Disease	Agro advisories	
Component	Condition			
General		• Punarvasu rainstar starts from July 6 th and remains upto July 19 th .		
		The normal rainfall of Punarvasu rainstar is 48.3 mm.		
		• Agro forestry on the bund involving <i>Melia dubia</i> (Hebbevu), Silver		
		oak and Causurina is recommended for sustainable income under		
		abnormal extreme rainfall events .		

		• Formore are advised to take up Disusking an accident the sold in
		 Farmers are advised to take up Ploughing or passing the cultivator in the areas where rainfall received and it should be taken up across the slope, it improves infiltration and conservation of rain water but also avoids soil erosion. The farmers are procure good quality of seeds and fertilizers. Take up the germination test for Kharif sowing. Take up land preparation and strengthen the bund with fodder grass which will facilitate conservation of rainwater due to pre-monsoon showers. Ploughing across the slope will make insitu soil moisture conservation
Vhanff	Couving	Time for application tank silt to increase soil fertility. The following energy are suggested for coming
Kharif crops	Sowing, Germination and Early vegetative stage	 The following crops are suggested for sowing. Under mono cropping the long duration crops: Redgram : TTB-7, BRG-1,2,4&5 Maize: Ganga-11, Deccan -103, Vijaya composite, Composite NAC-6004,6002, Hybrid-Nityashree (NAH-2049) Under double cropping the medium to short duration crops: Groundnut : TMV-2, JL-24, KCG-6 Cowpea: TVX-944, KBC-1, Green gram: PS-16, PDM-84,178 Black gram: Karagao-3, T-9, Rashmi(LBG-625). Redgram seeds treat with Carbendazim or Thiram @ 2 g/kg of seed 24 hours before sowing (or) with talc formulation of <i>Trichoderma viride</i> @ 4g/kg of seed (or) <i>Pseudomonas fluorescens</i> @ 10 g/kg seed and <i>Rhizobium</i> @ 10 g/kg of seed . Wherever, the long duration crops like Redgram, and Castor has been sown, undertake earthing up operation in addition to the above agronomic measures. Earthing up makes a better availability of soil moisture to the crop rows through conversion of the land into ridges and furrow system. Wherever over population of crop plants is existing, undertake thinning operation for better availability of scarce soil moisture to the reduced population of crop plants. Due to high humidity (warm and humid weather) possibility of pest and disease build up is more in already sown crop. Take prophylactic measures depending on the crop.
		Horticulture crop
Mango		 Deep ploughing is recommended in mango orchard. Rainwater harvesting through opening circular trenches around the tree at a distance of 6 feet and width at 9 inches, as well as depth and mulching the trenches with dry mango and green leaf manuring helps in retaining sufficient soil moisture in soil. Apply recommended dose of fertilizer in June month. Recommended for pruning of dried twigs and branches done in June-July month to facilitate the sunlight to bear the more fruits in next seasons.
		Animal Husbandry
Dairy		 Vaccinate the cow against the Black quarter disease. It is a bacterial disease caused by <i>Clostridium chauvoei</i> Feeding cow containing about 17 per cent dietary fiber in the animal feed are also helpful to increase fat percentage in milk. Concentrate mixture should comprise grains (40 per cent), oil cakes (32 per cent), brans (25per cent), mineral mixture (2 per

cent) and common salt (1 per cent).		
	Sericulture	
	1. Maintain the optimum room temperature in Sericulture unit.	
	1. Maintain the optimum room temperature in Serieutare and.	

AMFU of IMD Bengaluru

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*