

UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



Date: **07.10**.

AGROMET-ADVISORY BULLETIN

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

The forecast is valid for Chikkaballapur district Weather forecast (Valid from 08-10-2022 to 12-10-2022)

Forecast summary:

| Parameters | 08.10.2022 | 09.10.2022 | 10.10.2022 | 11.10.2022 | 12.10.2022 |
|---------------------------------|------------|------------|------------|------------|------------|
| Rainfall (mm) | 4 | 5 | 15 | 15 | 5 |
| Max Temp Trend (°C) | 28 | 27 | 28 | 28 | 28 |
| Min Temp Trend (°C) | 20 | 20 | 20 | 20 | 20 |
| Total cloud cover (octa) | 5 | 5 | 6 | 6 | 5 |
| Relative humidity (%)Max | 86 | 86 | 88 | 88 | 87 |
| Relative humidity (%)Min | 73 | 73 | 75 | 75 | 74 |
| Wind speed(Km/hr) | 5 | 6 | 5 | 4 | 4 |
| Wind Direction (Degrees) | 288 | 292 | 246 | 207 | 169 |

Light to moderate rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 27.0-28.0°C and Minimum of 20.0°C. Relative humidity 86-88 % during morning hrs and 73-75 % during noon is expected. Wind speed 4-6 km/hr.

Weather Based Agro Advisories Crop information and Crop Stages of the major Kharif/Rabi crops

| District | Kharif crops | | | Horticulture crops | | |
|----------------|--------------|---------|---------------|--------------------|-------|-------|
| Chikkaballapur | Groundnut | Redgram | Finger millet | Maize | Grape | Mango |
| a | PF,PM | V,F | V,T,E | Ts, CI,GF,FD | - | |

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

Agromet Advisory:

| Crop/ | Stage/ | Pest and Disease | Agro advisories | | |
|-----------|---------------|---|---|--|--|
| Component | Condition | | | | |
| General | | 1. Hasta rainstar starts from September 27 th and remains up to | | | |
| | | October 10 th . The normal rainfall of Hasta rainstar is 105.8 mm. | | | |
| Kharif | Vegetative | The following crops are suggested for sowing. | | | |
| crops | stage | Finger millet Ragi (Indaf-5, 7,9, ML-365, KMR-301), Horse gram- | | | |
| | And Tillering | PHG-9, KBH-1, Niger, Field bean-HA-3 and 4 and sunflower-KBSH- | | | |
| | stage | 1,41,42, 44 & 53, co | owpea (KBC-1, TVX-944 and PKB-4 for vegetable | | |
| | | purpose). | | | |
| | | Suggested for | or intercultivation for Ragi crop | | |
| | | | over population of crop plants is existing, undertake | | |
| | | 0 1 | eration for better availability of scarce soil moisture to population of crop plants. | | |
| | | | of leaf folder insect in Redgram to spray Methomyl | | |
| | | | m/litre of water before flowering of crop. | | |
| | | 4. If Wilt disea | ases noticed in Redgram field –Drenching Carbendazim itre of water. | | |

5. Remove and burn the infected Wilt diseases Redgram plant in field.

| | Horticulture crop | | | |
|-------|---|--|--|--|
| Mango | 1. Paclobutrazol soil application @ 5 g a.i per plant is recommended for uniform flowering behavior in Mango crop. Paclobutrazol is effective not only in flower induction but also in early and off season flower induction which thereby maintains regularity and synchronization in flowering. | | | |
| | Sericulture | | | |
| | Maintain the optimum room temperature in Sericulture unit. | | | |
| Dairy | Congenial climate and abundant fodder availability facilitate improved Dairy, Piggery, Poultry, Sheep and Goat rearing. | | | |
| | Feeding of lactating cow | | | |
| | Proper feeding of dairy cattle should envisage minimum wastage of nutrients and maximum returns in respect of milk produced. A concentrate mixture made up of protein supplements such as oil cakes, energy sources such as cereal grains (Maize, Jowar), tapioca chips and laxative feeds such as brans (rice bran, wheat bran, gram husk) is generally used. Mineral mixture containing major and all the trace elements should be included at a level of 2 percent. Dietary fiber for milking cow should be 17 % and NDF (Neutral Detergent Fiber) 22 % | | | |

AMFU of IMD Bengaluru

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*