

UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



Date: 25.08.2023

AGROMET-ADVISORY BULLETIN

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

The forecast is valid for Chikkaballapur district Weather forecast (Valid from 26-08-2023 to 30-08-2023)

Forecast summary:

Parameters	26.08.2023	27.08.2023	28.08.2023	29.08.2023	30.08.2023
Rainfall (mm)	0	1	1	1	1
Max Temp Trend (°C)	32	31	31	31	31
Min Temp Trend (°C)	20	20	20	20	20
Total cloud cover (octa)	5	5	5	5	5
Relative humidity (%) Max	79	79	79	79	79
Relative humidity (%) Min	71	71	71	71	71
Wind speed (Km/hr)	5	6	7	8	8
Wind Direction (Degrees)	288	291	292	291	292

Light rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 31.0-32.0°C and Minimum of 20.0°C. Relative humidity 79 % during morning hrs and 71 % during noon is expected. Wind speed 5-8 km/hr.

Weather Based Agro Advisories

Crop information and Crop Stages of the major Kharif/Rabi crops

District	Kharif crops			Horticulture crops		
Chikkaballapura	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
	F,PI	V	S,G,EV	V	-	-

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

Agromet Advisory:

Crop/ Component	Stage/ Condition	Pest and Disease	Agro advisories
General		 30th. The not One or two parts spells. Lack of soil recommender Mulching is 	tar starts from August 17 th and remains up to August rmal rainfall of Magha rainstar is 62.8 mm protective irrigation recommended due to prolonged dry moisture observed in crop field. Intercultural operation ed to conserve soil moisture and remove of weeds. recommended for conserve the soil moisture due to dry the Kharif crops
Kharif crops	Early vegetative, Vegetative stage and Flowering stage	Finger mille Cowpea: TV Black gram: Earthing up rows through Wherever ov thinning ope the reduce po	s are suggested for sowing. t: GPU-26,28,66 HR-911,KMR-204, ML-365 X-944, KBC-1, Green gram: PS-16, PDM-84,178 Karagao-3, T-9, Rashmi (LBG-625). makes better availability of soil moisture to the crop of conversion of the land into ridges and furrow system. Ver population of crop plants is existing, undertake tration for better availability of scarce soil moisture to opulation of crop plants. commended dose of Gypsum in Groundnut crop at the

	time of flowering
·	Horticulture crop
Mango	Deep ploughing is recommended in mango orchard. Rainwater harvesting through opening circular trenches around the tree at a distance of 6 feet and width at 9 inches, as well as depth and mulching the trenches with dry mango and green leaf manuring helps in retaining sufficient soil moisture in soil.
	Animal Husbandry
Dairy	 Vaccinate the cow against the Haemorrhagic Septecemia (HS) and Black Quarter (BQ) Disease at the age of six month old calf. Feeding of lactating cow Proper feeding of dairy cattle should envisage minimum wastage of nutrients and maximum returns in respect of milk produced. A concentrate mixture made up of protein supplements such as oil cakes, energy sources such as cereal grains (Maize, Jowar), tapioca chips and laxative feeds such as brans (rice bran, wheat bran, gram husk) is generally used. Mineral mixture containing major and all the trace elements should be included at a level of 2 percent. Dietary fiber for milking cow should be 17 % and NDF (Neutral Detergent Fiber) 22 %

AMFU of IMD Bengaluru

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*