

Forecast announce

UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



AGROMET-ADVISORY BULLETIN

Date: 28.07.2023

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department The forecast is valid for Chikkaballapur district Weather forecast (Valid from 29-07-2023 to 02-08-2023)

Forecast summary:					
Parameters	29.07.2023	30.07.2023	31.07.2023	01.08.2023	02.08.2023
Rainfall (mm)	3	1	2	2	1
Max Temp Trend (^o C)	29	31	31	31	31
Min Temp Trend (^o C)	22	22	22	22	22
Total cloud cover (octa)	5	5	5	5	5
Relative humidity (%) Max	85	85	83	83	83
Relative humidity (%) Min	74	74	72	72	72
Wind speed (Km/hr)	5	4	4	4	5
Wind Direction (Degrees)	292	292	291	291	288

Light to moderate rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 29.0-31.0°C and Minimum of 22.0°C. Relative humidity 83-85 % during morning hrs and 72-74 % during noon is expected. Wind speed 4-5 km/hr.

Weather Based Agro Advisories

Crop information and Crop Stages of the major Kharif/Rabi crops

District	Kharif crops			Horticulture crops		
Chikkaballapur	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
а	V,F	EV,V	S,G	EV,V	-	-

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

Agromet Advisory:

Crop/	Stage/	Pest and Disease	Agro advisories		
Component	Condition				
General		 Pushya rainstar starts from July 20th and remains up to August 02nd The normal rainfall of Pushya rainstar is 56.4 mm. Continue sowing operation due to sufficient soil moisture available in crop fields. Due to high humidity (warm and humid weather) possibility of pes 			
		prophylacticWind speed and HorticulWind speed	 and disease build up is more in already sown crop. Take prophylactic measures depending on the crop. Wind speed is high; postpone the spraying operation in Agriculture and Horticultural crops. Wind speed is high, provide staking support to Banana and vegetable crop to protect from uprooting of crops. 		
Kharif crops	Sowing, Germination and Early vegetative, Vegetative stage	 The following crops are suggested for sowing. Finger millet : MR-1,6, GPU-28, HR-911, ML-365 Cowpea: TVX-944, KBC-1, Green gram: PS-16, PDM-84,178 Black gram: Karagao-3, T-9, Rashmi(LBG-625). Earthing up makes a better availability of soil moisture to the cr rows through conversion of the land into ridges and furrow syst Wherever over population of crop plants is existing, undertake 			

	 thinning operation for better availability of scarce soil moisture to the reduced population of crop plants. Apply the recommended top dress fertilizer to Redgram and Maize crops. Horticulture crop
Mango	 Deep ploughing is recommended in mango orchard. Rainwater harvesting through opening circular trenches around the tree at a distance of 6 feet and width at 9 inches, as well as depth and mulching the trenches with dry mango and green leaf manuring helps in retaining sufficient soil moisture in soil.
	Animal Husbandry
Dairy	 Vaccinate the cow against the Black quarter disease. It is a bacterial disease caused by <i>Clostridium chauvoei</i> Feeding cow containing about 17 per cent dietary fiber in the
	2. Feeding cow containing about 17 per cent dictary fiber in the animal feed are also helpful to increase fat percentage in milk. Concentrate mixture should comprise grains (40 per cent), oil cakes (32 per cent), brans (25per cent), mineral mixture (2 per cent) and common salt (1 per cent).

AMFU of IMD Bengaluru

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*