

# UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



### AGROMET-ADVISORY BULLETIN

## Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

The forecast is valid for Chikkaballapur district Weather forecast (Valid from 01-11-2022 to 05-11-2022)

#### **Forecast summary:**

Parameters	01.11.2022	02.11.2022	03.11.2022	04.11.2022	05.11.2022
Rainfall (mm)	0	5	7	6	7
Max Temp Trend ( °C)	26	24	25	25	25
Min Temp Trend ( °C)	16	16	17	17	16
<b>Total cloud cover (octa)</b>	5	5	5	5	5
Relative humidity (%)Max	77	79	79	79	79
Relative humidity (%)Min	71	73	73	73	73
Wind speed(Km/hr)	2	3	4	4	5
Wind Direction (Degrees)	63	22	58	66	90

Light rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 24.0-26.0°C and Minimum of 16.0-17.0°C. Relative humidity 77-79 % during morning hrs and 71-73 % during noon is expected. Wind speed 2-5 km/hr.

# Weather Based Agro Advisories Crop information and Crop Stages of the major Kharif/Rabi crops

District	Kharif crops				Horticulture crops	
Chikkaballapur	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
a	PM,H	F	V,T,E,GF	GF,M,H	-	

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

### **Agromet Advisory:**

Crop/	Stage/ Condition	Pest and	Agro advisories		
Component		Disease			
General		1. Swathi rainstar starts from October 24 <sup>th</sup> and remains up to November 5 <sup>th</sup> . The normal rainfall of Swathi rainstar is 47.9 mm.			
Kharif	Tillering stage/Ear	The following crops are suggested for sowing.			
crops	head formation stage/flowering/pod formation and maturity stage	Finger millet Ragi (Indaf-5, 7,9, ML-365, KMR-301), Horse gram- PHG-9, KBH-1, Niger, Field bean-HA-3 and 4 and sunflower-KBSH-1,41,42, 44 & 53, cowpea (KBC-1, TVX-944 at PKB-4 for vegetable purpose).  1. Suggested for intercultivation for Ragi crop 2. To control of leaf folder insect in Redgram to specification of the second of the seco			

	Carbendazim 50 WP 2 g/litre of water.			
	5. Remove and burn the infected Wilt diseases Redgram plant in field.			
	6. To control Cob borer spray melathion 50 EC @ 2ml/litre (500			
	ml/acre)			
	Horticulture crop			
	Horticulture crop			
Mango	1. Paclobutrazol soil application @ 5 g a.i per plant is recommended for uniform flowering behavior in Mango crop. Paclobutrazol is effective not only in flower induction but also in early and off season flower induction which thereby maintains regularity and synchronization in flowering.			
	Sericulture			
	Maintain the optimum room temperature in Sericulture unit.			
Dairy	Congenial climate and abundant fodder availability facilitate improved Dairy, Piggery, Poultry, Sheep and Goat rearing.			
	Feeding of lactating cow			
	<ul> <li>Proper feeding of dairy cattle should envisage minimum wastage of nutrients and maximum returns in respect of milk produced.</li> <li>A concentrate mixture made up of protein supplements such as oil cakes, energy sources such as cereal grains (Maize, Jowar), tapioca chips and laxative feeds such as brans (rice bran, wheat bran, gram husk) is generally used.</li> <li>Mineral mixture containing major and all the trace elements should be included at a level of 2 percent.</li> </ul>			
	1. Dietary fiber for milking cow should be 17 % and NDF (Neutral Detergent Fiber) 22 %			

AMFU of IMD Bengaluru

**Important Note:** Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*