

UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



Date: 06.10.2023

AGROMET-ADVISORY BULLETIN

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

The forecast is valid for Bengaluru Rural district.

Weather forecast (Valid from 07-10-2023 to 11-10-2023)

Forecast summary:

Parameters	07.10.2023	08.10.2023	09.10.2023	10.10.2023	11.10.2023
Rainfall (mm)	1	0	3	4	4
Max Temp Trend (°C)	29	29	28	28	28
Min Temp Trend (°C)	20	20	20	20	20
Total cloud cover (octa)	5	5	5	5	5
Relative humidity (%)Max	79	79	79	81	81
Relative humidity (%)Min	61	61	61	63	63
Wind speed(Km/hr)	6	6	6	5	5
Wind Direction (Degrees)	292	30	117	112	90

Light rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 28.0-29.0°C and Minimum of 20.0°C. Relative Humidity 79-81 % during morning hrs and 61-63 % during noon is expected. Wind speed is 5-6 km/hr.

Weather Based Agro Advisories

Crop information and Crop Stages of the major Kharif/Rabi crops

District	Kharif crops				Horticulture crops	
Bangalore Rural	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
(BR)	PI,PF	F,PF	EV,V,E	Ts,CI,M	-	-

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching, CI: Cob Initiation

Agromet Advisory:

Crop/ Component	Stage/ Condition	Pest and Disease	Agro advisories	
General		 One or two protective irrigation to late sown Kharif crops is recommended Hasta rainstar starts from September 27th and remains up to October 10th. The normal rainfall of Hasta rainstar is 105.8 mm Intercultural operation recommended for conserving soil moisture and removal of weeds. Mulching is recommended to conserve the soil moisture in standing crops. 		
Kharif crops	Vegetative stage and Flowering stage	The following crops are suggested for sowing. Finger millet Ragi (Indaf-5, 7,9, ML-365, KMR-301), Horse gram-PHG-9, KBH-1, Niger, Field bean-HA-3 and 4 and sunflower-KBSH-1,41,42, 44 & 53, cowpea (KBC-1, TVX-944 and PKB-4 for vegetable purpose). • Earthing up makes better availability of soil moisture to the crop rows through conversion of the land into ridges and furrow system. • Wherever over population of crop plants is existing, undertake		

	thinning operation for better availability of scarce soil moisture to			
	the reduce population of crop plants.			
	• To control leaf folder insect in Redgram spray Methomyl 40SP, 2			
	gram/litre of water before flowering of crop.			
	Wilt disease noticed: Redgram field-Drench with Carbendazim 50			
	WP 2 g/litre of water.			
	• For Biological control of pod borer insect in Redgram 200 LE/acre			
	HANPV 400 litre with teepol (1 ml/lit.). Pheromone traps			
	for Helicoverpa armigera 12/ha			
	 Remove and burn the infected Wilt diseased Redgram plant in field. 			
	Animal Husbandry			
Dairy	Feeding of lactating cow			
	 Proper feeding of dairy cattle should envisage minimum wastage of nutrients and maximum returns in respect of milk produced. 			
	• A concentrate mixture made up of protein supplements such as oil			
	cakes, energy sources such as cereal grains (Maize, Jowar), tapioca			
	chips and laxative feeds such as brans (rice bran, wheat bran, gram			
	husk) is generally used.			
	Mineral mixture containing major and all the trace elements should			
	be included at a level of 2 percent.			
	•			
	Detergent Fiber) 22 %			
	• Dietary fiber for milking cow should be 17 % and NDF (Neutral			

AMFU of IMD Bengaluru

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*