

UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



Date: 24.05.2022

AGROMET-ADVISORY BULLETIN

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

The forecast is valid for Bengaluru Rural district Weather forecast (Valid from 25-05-2022 to 29-05-2022)

Forecast summary:

Parameters	25.5.2022	26.5.2022	27.5.2022	28.5.2022	29.5.2022
Rainfall (mm)	1	2	0	0	0
Max Temp Trend (°C)	32	32	30	32	32
Min Temp Trend (°C)	21	21	20	20	21
Total cloud cover (octa)	6	7	7	4	4
Relative humidity (%)Max	82	82	87	87	85
Relative humidity (%)Min	55	55	58	57	55
Wind speed(Km/hr)	6	6	7	6	6
Wind Direction (Degrees)	270	270	249	252	252

Light rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 30.0-32.0°C and Minimum of 20.0-21.0°C. Relative 82-87 % during morning hrs and 55-58 % during noon is expected. Wind speed is 6-7 km/hr.

Weather Based Agro Advisories

Crop information and Crop Stages of the major Kharif/Rabi crops

District	Kharif crops Horticulture crops				ture crops	
Bangalore Rural	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
(BR)					-	FD,M

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching, CI: Cob Initiation

Agromet Advisory:

Crop/	Stage/	Pest and Disease	Agro advisories
Component	Condition		
General		over some Southeast A central Bay during next Rohini rains normal rainf The farmers up the germ Take up lan which will the showers. Ple conservation Taken up s	star starts from May 25 th and remains upto June 7 th . The fall of Rohini rainstar is 61.9 mm. Is are procure good quality of seeds and fertilizers. Take ination test for Kharif sowing. It is depreparation and strengthen the bund with fodder grass facilitate conservation of rainwater due to pre-monsoon oughing across the slope will make insitu soil moisture

		 Time for application tank silt to increase soil fertility. Sufficient rainfall received in May month sowing of long duration crops like Redgram BRG-1, 2, 4 &5, TTB-7 and Castor DCH-9, 32 &177 is recommended. 			
	Horticulture crop				
Mango	Fruit Development and Maturity stage	 In mango, fruit fly damage is also noticed. suitable control measure by spraying Carbaryl 4 g/litre of water Harvesting of mango without dropping ground level using mango net. Mango growers should harvest matured fruit and try to leave one or two inches of stem in the mango bunch, which helps to avoid sap burn disease. Afterward, the mango fruit should be wrapped by using papers and then keep it in plastic crates. After within 24 hours of harvesting, the crates are placed on top of each other with 5 to 6 create half-feet apart and then take 300 ml. of water in small plastic bottles, add 10 ml of ethylene to water, then after adding 10 g sodium hydroxide (caustic soda). Then seal the tarpaulin cover tightly without aeration for 48 to 60 hours, then tarpaulin should be removed and left out for aeration. Ethylene should be kept in 8 to 10 glasses for to ripen 1 ton of fruit. Natural ripening will help to get good quality fruit and better market price. 			
		Sericulture			
	1. Maintain	the optimum room temperature in Sericulture unit.			

AMFU of IMD Bengaluru

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*