

UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU GRAMIN KRISHI MAUSAM SEWA(GKMS) AMFU OF IMD, BENGALURU



Date: 28.11.2023

AGROMET-ADVISORY BULLETIN

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

The forecast is valid for Chikkaballapur district Weather forecast (Valid from 29-11-2023 to 03-12-2023)

Forecast summary:

Parameters	29.11.2023	30.11.2023	01.12.2023	02.12.2023	03.12.2023
Rainfall (mm)	0	1	2	2	0
Max Temp Trend (°C)	26	27	27	27	27
Min Temp Trend (°C)	18	17	17	18	18
Total cloud cover (octa)	5	6	7	8	8
Relative humidity (%) Max	89	89	91	91	91
Relative humidity (%) Min	76	76	78	78	78
Wind speed (Km/hr)	4	4	3	5	4
Wind Direction (Degrees)	112	103	69	58	11

Light rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 26.0-27.0°C and Minimum of 17.0-18.0°C. Relative humidity 89-91 % during morning hrs and 76-78 % during noon is expected. Wind speed 3-5 km/hr.

Weather Based Agro Advisories

Crop information and Crop Stages of the major Kharif/Rabi crops

District	Kharif crops			Horticulture crops		
Chikkaballapura	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
	-	М,Н	М,Н	-		-

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

Agromet Advisory:

Crop/ Component	Stage/ Condition	Pest and Disease	Agro advisories	
General		• Intercultural operation recommended for conserving soil moisture and removal of weeds.		
		• Mulching is reco crops.	mmended to conserve the soil moisture in standing rabi	
Kharif	Maturity and	The following Rabi crops are suggested for sowing.		
crops	Harvesting	Horse gram-PHG-9	O, KBH-1, Niger, Field bean-HA-3 and 4 and sunflower-	
	stage	KBSH-1, 41, 42, 44 & 53, cowpea (KBC-1, TVX-944 and PKB-4 for		
		vegetable purpose).		
		• Earthing up makes better availability of soil moisture to the crop rows through conversion of the land into ridges and furrow system.		
		thinning ope	ver population of crop plants is existing, undertake eration for better availability of scarce soil moisture to opulation of crop plants.	
		• Right time for	or harvesting of matured pigeon pea pod or whole plants	
		kept for dry	place.	
		Animal	Husbandry	

Dairy	Feeding of lactating cow
	• A concentrate mixture made up of protein supplements such as oil
	cakes, energy sources such as cereal grains (Maize, Jowar), tapioca
	chips and laxative feeds such as brans (rice bran, wheat bran, gram
	husk) is generally used.
	 Mineral mixture containing major and all the trace elements should
	be included at a level of 2 percent.
	• Dietary fiber for milking cow should be 17 % and NDF (Neutral
	Detergent Fiber) 22 %

AMFU of IMD Bengaluru

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: *mausam.imd.gov.in*