

**UNIVERSITY OF AGRICULTURAL SCIENCES, BENGALURU &
INDIAN METEOROLOGICAL DEPARTMENT**



**GRAMIN KRISHI MAUSAM SEWA
AMFU, OFRS, NAGANAHALLI,
MYSURU - 570003**



Date: 02-04-2024

AGRO-ADVISORY BULLETIN FOR MANDYA DISTRICT

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

Past Weather Data			
Parameter	30.03.2024	31.03.2024	01.04.2024
Rainfall (mm)	0	0	0
Max. Temp. (°C)	37.6	37.2	36.6
Min. Temp. (°C)	19.9	20.3	21.8
Sky condition (Octas)	2	2	4
Relative humidity (%) 0830 hours	65	77	70
Relative humidity (%) 1730 hours	30	33	33
Wind Speed (km/h)	0	4	4
Wind Direction	0	180	230

Weather forecast for the next five days (From 03-03-2024 to 07-04-2024)

Parameter	03.04.2024	04.04.2024	05.04.2024	06.04.2024	07.04.2024
Rainfall (mm)	0	0	0	0	4
Max. temp (°C)	37	37	37.5	38.1	37
Min.Temp (°C)	19	19	20	20	21
Sky condition (Octas)	2	2	4	4	4
Relative humidity (%) 0830 hours	76	74	75	58	80
Relative humidity (%) 1730 hours	20	23	19	17	15
Wind Speed (kmph)	9	9	10	10	14
Wind Direction	225	245	250	190	115

Forecast Summary

As forecast received from IMD, Partially cloudy sky with very light rainfall may be expected from 03.04.2024 to 07.04.2024 in Mandya district. The day temperature is expected to be 37-38.1°C & night temperature is expected 19-21°C. The relative humidity in the morning hours is expected to be 58-80% & afternoon relative humidity is expected to be in the range of 15-23% per cent. Wind speed expected to be 9-14 km/ hr.

Recommendations to the farmers:

Crop	Pest/Disease	Damage symptoms	Control measures
<ul style="list-style-type: none"> Mulch the crop with crop residues it will minimize water loss through evapotranspiration. Farmers should provide protective irrigation if rainfall does not occur at the critical stage of crop growth. Take deep summer ploughing after harvesting of Rabi crops, due to overturning of the soil in summer ploughing the sharp rays of sun enters the soil and kills the eggs, larvae and pupae of pests. This avoids the spread of the pests in the next growing season. Due to dry weather prevail for next four days, farmers may take up irrigation to the dry land horticultural crops and other field crops depending on soil and crop condition. As dry weather is predicted for the next five days, care should be taken to maintain the optimum moisture level of 60% in vermicompost pits. 			
Chilli	Thrips and mites	Symptoms : Severe leaf curling in both upward and downward directions, elongated petiole, Yellow mixed green leaves, the irregular leaf size with curled leaves, stunted growth of the plants.	Management: spray 2.5 ml. Dicofol 18.5 EC or 0.3 ml. Imidacloprid 17.8 SL./lit. water. Repeat the spray at an interval of 10 days, around 200 lit. spray solution/acre/spray.
Bean	Yellow mosaic virus	Yellow mixed green coloured leaves, stunted growth, to manage this disease uproot the infected plants and spray Imidocloprid 17.8 SL @ 0.5ml/lit.of water.	
Maize	Fall army worm	5% damage- Application of 5% neem seed kernel extract (NSKE) or azadirachtin 1500ppm @ 5ml/litre (1 litre/acre)of water. 10% damage- Whorl application of any one of the recommended insecticides for FAW, viz.,Chlorantraniliprole 18.5 SC (80 ml/acre) @ 0.4 ml/litre or Spinetoram 11.7 %SC (100ml/acre) @ 0.5 ml/litre or Emamectin benzoate 5% SG (80g/acre) @ 0.4g/litre.	
Banana	Pseudo stem borer	<ul style="list-style-type: none"> The grubs bore the pseudostem, affected plants with number of holes on pseudostem, breaking of infested plants at damaged portion due to wind. Management: Add 3.0 ml. Chloropyriphos 20 EC.or 3.0 ml. Dimethoate 30 EC. with 5.0 ml. water. Inject the mixture to the pseudostem about a feet height from ground level. 	
Tomato	Fruit borer	For every 25 rows of tomato grow one row of marigold cultivar African tall. The marigold seedlings about 35-40 days old. If borer problems exceeds 10 per cent spray 4 per cent. NSKE or 1.0 g. Methomil 40 SP. in a lit. water.	
Cowpea/Black gram	Aphids	Several individuals suck the sap from tender shoots, flower buds, flowers, leaves and tender pods. Yellowing of leaves, shrivelled grains and sooty mold on leaves. To manage this pest spray Dimethoate – 30 EC. @ 1.7 ml./liter water spray. 200 lit. Spray	

		mixture/acre or azadirachtin 1500ppm @ 5ml/litre.
watermelon	Sucking pests	For control of sucking insects in water melon install 8 – 10 yellow sticky traps per acre. When the infestation is sever spray melathion 50 EC @ 1 ml/lit water.
Cattle management during summer months	<ul style="list-style-type: none"> • Maximum temperature expected to rise in coming days so tie the livestock animals in shade / cool place. • Put 4-6 inches of grass cover on the cattle shed, water 2-3 times in the afternoon and also water the floor and walls of the cattle shed. This keeps the cattle shed fairly cool. • Arrangements should be made so that air can flow easily in the cattle shed. • Clean water should be given to the cattles. • Don't allowed the cattle to graze during afternoon period when the temperature is high, cattles should be allowed to graze in the morning or evening time. 	

Block level weather forecast (From 03-07-2024 to 07-04-2024)

Krishnarajpet					
Parameter	03.04.2024	04.04.2024	05.04.2024	06.04.2024	07.04.2024
Rainfall (mm)	0	0	0	0	0.7
Max. temp (°C)	36.1	35.8	35.7	36.4	36.8
Min.Temp (°C)	18.8	18.8	19.1	19.1	18.9
Sky condition (Octas)	2	2	3	4	2
Relative humidity (%) 0830 hours	84	76	73	76	76
Relative humidity (%) 1730 hours	17	22	21	19	18
Wind Speed (kmph)	9	10	10	8	11
Wind Direction	251	251	251	248	248

Maddur					
Parameter	03.04.2024	04.04.2024	05.04.2024	06.04.2024	07.04.2024
Rainfall (mm)	0	0	0	0	0
Max. temp (°C)	37.4	37.3	36.9	37.4	38.2
Min.Temp (°C)	22.2	21.7	21.9	21.7	21.8
Sky condition (Octas)	3	2	2	4	1
Relative humidity (%) 0830 hours	82	83	81	87	86
Relative humidity (%) 1730 hours	15	18	18	18	15
Wind Speed (kmph)	9	10	8	8	12
Wind Direction	248	243	68	214	117

Malvalli

Parameter	03.04.2024	04.04.2024	05.04.2024	06.04.2024	07.04.2024
Rainfall (mm)	0	0	0	0	0
Max. temp (°C)	36	36.2	36.2	36.4	37.2
Min.Temp (°C)	21.5	20.7	21.1	20.9	20.8
Sky condition (Octas)	3	1	2	3	1
Relative humidity (%) 0830 hours	80	86	80	85	84
Relative humidity (%) 1730 hours	16	18	17	16	15
Wind Speed (kmph)	9	11	10	9	12
Wind Direction	248	248	252	248	248

Mandya

Parameter	03.04.2024	04.04.2024	05.04.2024	06.04.2024	07.04.2024
Rainfall (mm)	0	0	0	0	0
Max. temp (°C)	36.5	36.4	36.2	36.7	37.7
Min.Temp (°C)	20.9	20.5	21	20.7	20.7
Sky condition (Octas)	3	2	2	4	1
Relative humidity (%) 0830 hours	84	81	82	89	85
Relative humidity (%) 1730 hours	15	19	18	18	15
Wind Speed (kmph)	11	11	10	10	13
Wind Direction	248	248	248	244	248

Nagamangala

Parameter	03.04.2024	04.04.2024	05.04.2024	06.04.2024	07.04.2024
Rainfall (mm)	0	0	0	0	0.8
Max. temp (°C)	37.3	36.6	36.3	37.1	37.7
Min.Temp (°C)	20.7	20.8	20.6	20.9	20.8
Sky condition (Octas)	2	2	3	4	2
Relative humidity (%) 0830 hours	80	74	75	76	76
Relative humidity (%) 1730 hours	14	21	21	19	18
Wind Speed (kmph)	11	10	10	9	12
Wind Direction	248	117	243	68	117

Pandavapura



Parameter	03.04.2024	04.04.2024	05.04.2024	06.04.2024	07.04.2024
Rainfall (mm)	0	0	0	0	0.2
Max. temp (°C)	35.9	35.6	35.5	36	36.9
Min.Temp (°C)	19.3	19.4	19.9	19.6	19.6
Sky condition (Octas)	3	2	3	4	1
Relative humidity (%) 0830 hours	84	76	76	83	83
Relative humidity (%) 1730 hours	18	22	20	19	17
Wind Speed (kmph)	10	10	9	10	13
Wind Direction	252	252	248	248	248

Shrirangapattana

Parameter	03.04.2024	04.04.2024	05.04.2024	06.04.2024	07.04.2024
Rainfall (mm)	0	0	0	0	0.1
Max. temp (°C)	35.1	34.9	34.8	35.3	36.3
Min.Temp (°C)	19.1	19	19.6	19.2	19.2
Sky condition (Octas)	3	1	3	4	1
Relative humidity (%) 0830 hours	85	78	76	84	84
Relative humidity (%) 1730 hours	18	22	19	19	17
Wind Speed (kmph)	10	11	10	10	13
Wind Direction	252	248	244	248	248


NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 11
(12.03.2024 to 18.03.2024) over Agricultural region

Karnataka




Agriculture vigour is good at isolated places of coastal Karnataka.


Vegetation Condition Index



Temperature Condition Index



Forecasted soil moisture



Farmers facing high temperatures, low soil moisture and reduced vegetation should prioritize efficient irrigation techniques, such as drip irrigation and implement mulching to conserve moisture and mitigate evaporation. Opting for drought-resistant crops and employing soil conservation methods like cover cropping can help maintain soil health and resilience.

- Download “**DAMINI**” app to get early warning on lightening and take precautions based on the alert given by the application.
- Kindly download “**MAUSAM**” APP for location specific forecast & warning & “**MEGHDOOT**” APP for Agromet advisory
- This information is available in the website: mausam.imd.gov.in

For any information farmers can contact **Dr.C.Ramachandra**, Senior Farm Superintendent/ **Dr. Sumanth Kumar.G.V**, Research Assocateover phone No. 0821-259126/ 9535345814.

AMFU of IMD,
Naganahalli, Mysuru