

**UNIVERSITY OF AGRICULTURAL SCIENCES, BENGALURU &
INDIAN METEOROLOGICAL DEPARTMENT**



**GRAMIN KRISHI MAUSAM SEWA
AMFU, OFRS, NAGANAHALLI,
MYSURU - 570003**



Date: 14-06-2024

AGRO-ADVISORY BULLETIN FOR KODAGU DISTRICT

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

Past Weather Data			
Parameter	12.03.2024	13.03.2024	14.03.2024
Rainfall (mm)	12.5	0	-
Max. Temp. (°C)	26.4	26	-
Min. Temp. (°C)	21.2	20	-
Sky condition (Octas)	-	-	-
Relative humidity (%) 0830 hours	100	99	-
Relative humidity (%) 1730 hours	97	94	-
Wind Speed (km/h)	-	-	-
Wind Direction	-	-	-

Weather forecast for the next five days (From 15-06-2024 to 19-06-2024)

Parameter	15.06.2024	16.06.2024	17.06.2024	18.06.2024	19.06.2024
Rainfall (mm)	5	3	10	12	15
Max. Temp. (°C)	27	28	27	27	26
Min. Temp. (°C)	17.6	17.4	18.2	18.6	17.6
Sky condition (Octas)	7	7	7	8	8
Relative humidity (%) 0830 hours	94	93	92	94	95
Relative humidity (%) 1730 hours	62	53	60	60	86
Wind Speed (kmph)	9	8	9	9	8
Wind Direction	236	238	225	212	202

Forecast Summary

As forecast received from IMD, cloudy sky with **light to moderate rainfall** may be expected from 15.06.2024 to 19.06.2024 in Kodagu district. The day temperature is expected to be 26-28°C & night temperature is expected 17.4-18.6°C. The relative humidity in the morning hours is expected to be 92-95% & afternoon relative humidity is expected to be in the range of 53-86%. Wind speed expected to be 8-9 km/ hr.

Recommendations to the farmers:

Crop	Pest/Disease	Damage symptoms	Control measures
Crops that can grown in the month of June			
Paddy: BR-2655, Jaya, MSN-99			
Aerobic Rice: Daksha (KMP-175), Vanasiri (MAS-26), Sharada (MAS-946), Anaga (ARB-6), Rashi			
Finger millet: INDAF-8, MR-1, MR-6, L-5, KMR-301, GPU-45, KMAR-316			
Kodo millet: RBK-155, GPUK-3			
Maize: Hema, Nityashree, MAH-14-5			
Shaktiman Maize: CSH-5, CSH-9, CSV-4			
Pop corn: Amber			
Little millet: Co-2, OLM-203, GPUL-6			
Barugu: GPUP-8, GPUP-21, GPUP-28			
Navane: PS-4, SIA-326, GPUF-3			
Barnyard millet: DHBM-93-3			
Bajra: WCC-75, PHVB-910, ICTP-8203, SMV-221, MH-946			
Groundnut: GKVK-5, GKVK-27, KCG-6, GpBD-4, ICGV,-91114, JL-24, TMV-2			
Sunflower: KBSH-41, KBSH-42, KBSH-44, KBSH-53, KBSH-78, KBSH-85			
Castor: DSH-9, DSH-177, GSH-4, 48-1			
Niger: KBN-1			
Red gram: BRG-1, BRG-2, BRG-36, BRG-4, BRG-5, ICP-7035, TTB-7, Hyderabad-3C			
Sugarcane: Co-62175, Co-86032, CoVC-18061			
Horticultural crops: Tomato, Potato, Brinjal, Chilli, Cabbage, Radish, Knol Khol, Carrot, Onion, Mango, Banana, Lemon, Guava, Mosambi, Sapota, papaya, Pomegranate, Pineapple, Cashew nut, Garlic, Ginger, Tumeric			
Fodder crops:			
Maize: African tall			
Sorghum: MP Chaari, Pusa Chaari, JS-3, JS-20, COFS-29			
Bajra: Deena Bhandu-49A			
Cowpea: KBC-2, MFC-09-3			
Water Management:			
✓ Irrigation: Suspend irrigation activities to prevent overwatering.			
✓ Drainage: Ensure proper drainage systems are in place to avoid waterlogging in the fields.			
Rainwater Harvesting: Set up rainwater harvesting systems to capture and store rainfall for future use.			
Seed treatment of Green gram, black gram, cowpea, Pigeon pea			
Treat the seeds with bio-fertilizers like Rhizobium and Phosphorus Solubilising Bacteria (PSB) @ 200 g. /acre, each.			
Cowpea	Aphids	Dimethoate - 30 EC. @ 1.7 ml./liter water spray. 200 lit. spray mixture/acre	
Coconut	Rugose whitefly	<ul style="list-style-type: none"> ✓ Spraying starch solution (1%) to dislodge the heavy sooty mould deposition on the leaves of infested plants. ✓ Use of yellow sticky traps to trap the adult whiteflies ✓ In case of severe infestation, spray neem oil 0.5% 	
Black gram	Aphid	Adults and nymphs suck the sap from tender shoots, leaves; flower buds, flowers and tender pods. Yellowing of leaves,	

		shrivelled grain and development of sooty mold. Spray 1.7 ml. Dimethoate 30 EC. in lit. water, when infestation noticed. 250 lit. spray solution/acre.
Cowpea	Pod borer	Caterpillar bore the pods and feed on seeds. Infested pods with hole. On tender pods spray Chlorpyriphos - 20 EC. @ 2 ml./lit. water. 300 lit. spray solution/acre.
Sugarcane	Seed treatment	Treat the sets with 1 g. Carbendazim/ liter of water for 15 minutes followed by transplanting
Chilli	Use of growth regulator	<ul style="list-style-type: none"> • Generally flower drop will be noticed in chilli crop. • To prevent flower drop spray NAA (Planofix) 50 ppm (50 mg. in 1 litre water) at time of flowering results in higher yield.
Tomato	Serpentine leaf miner	Larva mines the leaf epidermis, serpentine tunnels on leaves. Spray 0.3 ml. Imidachloprid 17.8 SL. or 2.0 ml. Triazophos 40 EC/lit. water.
Tomato	Fruit borer	<ul style="list-style-type: none"> • Caterpillar bore the flower buds and fruits. Infested flower buds with hole and drops off, fruit with a hole, water enter through the hole leads to rotting. • Trap crop: For every 25 rows of tomato grow one row of marigold cultivar African tall. The marigold seedlings about 35-40 days old. If borer problems exceeds 10 per cent spray 4 per cent. NSKE or 100 LE, Ha. NPV. If infestation in severe form spray 1.0 g. Methomil 40 SP. in a lit. water
Coconut	Rhinoceros beetle	Remove the beetle from infected part and fill 2 % Quinolphos or 5 % Melathion in sand @ 1:1 ratio.
Ginger	Rhizome treatment	Treat the rhizomes with 2 gm bleaching powder, 1 gm Metalaxyl MZ- 72, 1 gm Streptocycline mixed in 1 ltr of water. Treat the rhizomes 30 minutes and dry in shade. This will prevent rhizome rot problems.
Ginger	Rhizome rot	<p>Symptoms: Yellowing of leaves, drying of leaf margin followed by burning of entire leaf, bottom portion of the shoot with water soaked soft, orange coloured rhizomes turn to brown colour, gradual rotting of such rhizomes.</p> <p>To manage this disease by drench the base of the plants with 2.0 g. Captan 50 WP or 2.0 g. Metalaxyl - MZ 72WP. In a lit.water.</p>
Sugarcane	Topdressing and intercultivation	Provide topdressing in 3 splits, first topdressing should be given in 45 days after planting with 20% nitrogen, second is to be given after 75 days after planting using 30% of total nitrogen. The final topdressing should be given after 105 days with the remaining 40% nitrogen. Application of fertilizers in between ploughed rows of both sides will be more efficient use of fertilizers possible. Alternatively provide the fertilizers surrounding seedlings with 4 - 5" depth.

Block level weather forecast (From 15-06-2024 to 19-06-2024)**Madikeri**

Parameter	15.06.2024	16.06.2024	17.06.2024	18.06.2024	19.06.2024
Rainfall (mm)	9.5	25.9	2	3.3	3.6
Max. temp (°C)	26.6	24.8	26	30.1	29.9
Min.Temp (°C)	20.8	20.2	19.6	20.5	20.5
Sky condition (Octas)	8	8	8	8	8
Relative humidity (%) 0830 hours	97	98	98	98	98
Relative humidity (%) 1730 hours	84	94	88	72	73
Wind Speed (kmph)	8	8	7	7	7
Wind Direction	283	248	249	257	248

Somvarpet

Parameter	15.06.2024	16.06.2024	17.06.2024	18.06.2024	19.06.2024
Rainfall (mm)	5.3	13.8	0.5	0	0
Max. temp (°C)	24.8	23.8	25.4	29.3	29.4
Min.Temp (°C)	19.3	18.6	18.5	18.9	19
Sky condition (Octas)	8	8	8	8	8
Relative humidity (%) 0830 hours	94	95	96	92	96
Relative humidity (%) 1730 hours	82	88	80	63	64
Wind Speed (kmph)	14	15	14	13	12
Wind Direction	252	248	249	257	249

Virajpet

Parameter	15.06.2024	16.06.2024	17.06.2024	18.06.2024	19.06.2024
Rainfall (mm)	5.9	23.1	1.5	3.6	3.4
Max. temp (°C)	28	24.3	25.5	30.3	29.7
Min.Temp (°C)	21.6	21	20.3	21.6	21.2
Sky condition (Octas)	8	8	8	8	8
Relative humidity (%) 0830 hours	96	97	98	97	97
Relative humidity (%) 1730 hours	75	93	87	70	70
Wind Speed (kmph)	7	7	6	7	7
Wind Direction	248	248	248	248	248

- Download “**DAMINI**” app to get early warning on lightening and take precautions based on the alert given by the application.
- Kindly download “**MAUSAM**” APP for location specific forecast & warning & “**MEGHDOOT**” APP for Agromet advisory
- This information is available in the website: mausam.imd.gov.in

For any information farmers can contact Dr. C. Ramachandra, Senior Farm Superintendent/ Dr. Sumanth Kumar.G.V, Research Assoicate over phone No. 0821-259126/ 9535345814.

AMFU of IMD,
Naganahalli, Mysuru