



UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU
GRAMIN KRISHI MAUSAM SEWA(GKMS)
AMFU OF IMD, BENGALURU



AGROMET-ADVISORY BULLETIN

Date: **09.12.2022**

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

**The forecast is valid for Chikkaballapur district
Weather forecast (Valid from 10-12-2022 to 14-12-2022)**

Forecast summary:

Parameters	10.12.2022	11.12.2022	12.12.2022	13.12.2022	14.12.2022
Rainfall (mm)	1	17	11	14	14
Max Temp Trend (°C)	22	22	25	26	27
Min Temp Trend (°C)	19	19	19	19	19
Total cloud cover (octa)	5	7	7	7	5
Relative humidity (%)Max	57	64	64	64	60
Relative humidity (%)Min	44	50	50	50	45
Wind speed(Km/hr)	3	4	5	4	4
Wind Direction (Degrees)	66	22	27	112	112

Light rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 22.0-27.0°C and Minimum of 19.0°C. Relative humidity 57-64 % during morning hrs and 44-50 % during noon is expected. Wind speed 3-4 km/hr.

**Weather Based Agro Advisories
Crop information and Crop Stages of the major Kharif/Rabi crops**

District	Kharif crops				Horticulture crops	
Chikkaballapur	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
a	--	PF,GF,M	M,H	--	-	--

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Transplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tilling, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

Agromet Advisory:

Crop/Component	Stage/Condition	Pest and Disease	Agro advisories
General			<ul style="list-style-type: none"> Cyclonic Storm “Mandous” pronounced as “Man-Dous” formation over Southwest Bay of Bengal resulted in cloudy weather and light to moderate rainfall forecasted from IMD. Jyestha rainstar starts from December 3rd to December 15th. The normal rainfall of Jyestha rainstar is 7.7 mm. The following crops are suggested for sowing. Field bean-HA-3 and 4 and sunflower-KBSH-1,41,42, 44 & 53, cowpea (KBC-1, TVX-944 and PKB-4 for vegetable purpose). Horse gram- PHG-9, KBH-1 The grains of the harvested crops should be properly dried by retaining moisture percentage of Cereals 11-12 %, Pulses-9%, Oilseeds-8% and Vegetable seeds 5-6% for long storage & also minimize the store pest damage. To protect the pulse grains from storage pests apply oils of Castor/ linseed/honge/neem oil @ 3-5 ml per kg of grains.

Finger millet	Harvesting	1. Advised for postpone harvesting next couple of days due to cloudy and light rainfall forecasted from IMD
Maize	Harvesting	1. Advised for postpone harvesting next couple of days due to cloudy and light rainfall forecasted from IMD
Horticulture crop		
Mango	Flower bud initiation and Flowering	<ol style="list-style-type: none"> 1. Clear the weeds in Mango orchard/Guava/Sapota put it under the basin as mulch. 2. If Phanerogamic plants are growing on the mango tree to cut/prune out completely and apply Bordeaux paste or Copper oxy chloride that portion. 3. Remove the weeds such as lantana which are growing under the mango tree. 4. Leaf hopper and Powdery mildew disease incidence is more before flowering and immediately after fruit formation to manage spraying of Carbaryl, 50WP @4g/litre of water or Imidachlorprid @ 0.3ml/ litre of water for management of leaf hopper. 5. Spray Lamda Cyhalothrin 5EC @ 0.5 ml/ litre of water or sulphur dust (SULTAF) 80 W @3g/litre of water against the Powdery mildew diseases. 6. If the incidence of Leaf hopper is severe spray Azadirachtin (10,000 ppm) @ 7.0 ml/ litre of water.
Animal Husbandry		
		<ol style="list-style-type: none"> 1. To protect animals from a sudden drop in temperature, keep the animals in a covered shed/area during the night. The bedding/hay in the animal sheds must be kept dry and changed/aired every day. 2. Due care should be taken to store/procure fodder for periods of shortage that may occur during the winter months in certain areas. Perennial grasses must be cut at this time.
Sericulture		
		<ol style="list-style-type: none"> 1. White muscardine: caused by <i>Beauveria bassiana</i> , 2. Manage the humidity in the rearing house by providing good cross-ventilation. Dust dry slaked lime powder when silkworms settle for moult. 3. Feed silkworms with adequate quantity of mulberry leaves to avoid the accumulation of left over leaves in the rearing bed. Make sure that the silkworm bed is dry and thin. 4. If the silkworm rearing house temperature falls below 22°C, raise it using room heater / charcoal stove. 5. Collect muscardine affected larvae from the rearing bed before mummification, dust antimuscardine bed disinfectant and finally burn them. Do not throw them on the street or feed to animals / birds. 6. Dust Vijetha and Vijetha Supplement or Ankush bed disinfectant as per recommended schedule or dust any recommended anti-muscardine bed disinfectant as per the schedule.
Poultry		
		<ol style="list-style-type: none"> 1. The poultry house should be located in such a way that long axis is in east-west direction. This will prevent the direct sunshine over the birds. 2. Beginning at one day of age, the chick should be housed at a temperature 35° C will maintain one week, at a relative humidity between 40 – 60% after wards 2 to 4 weeks temperature decreases every week by 2° C. 3. Provide artificial brooding to chicks to maintain adequate temperature. 4. Care should be taken to prevent the chicks from being exposed to wind chill.

- | | |
|--|--|
| | <ol style="list-style-type: none">5. Sides should be covered with curtains during cool hours of the day.6. Wet litter material should be removed regularly7. Ensure proper cross ventilation to avoid ammonia accumulation |
|--|--|

**AMFU of IMD
Bengaluru**

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website:
mausam.imd.gov.in