

**UNIVERSITY OF AGRICULTURAL SCIENCES, BENGALURU &  
INDIAN METEOROLOGICAL DEPARTMENT**



**GRAMIN KRISHI MAUSAM SEWA  
AMFU, OFRS, NAGANAHALLI,  
MYSURU - 570003**



Date: 21-06-2024

**AGRO-ADVISORY BULLETIN FOR CHAMARAJANAGARA DISTRICT**  
Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

**Weather forecast for the next five days (From 22-06-2024 to 26-06-2024)**

Parameter	22.06.2024	23.06.2024	24.06.2024	25.06.2024	26.06.2024
Rainfall (mm)	3	8	9	5	6
Max. Temp. (°C)	31	30	29	28	28
Min.Temp. (°C)	16.7	16.4	17.9	18	17.7
Sky condition (Octas)	8	8	8	8	8
Relative humidity (%) 0830 hours	79	85	86	83	86
Relative humidity (%) 1730 hours	49	60	63	68	63
Wind Speed (kmph)	18	13	16	18	21
Wind Direction	252	248	248	248	248

**Forecast Summary**

As forecast received from IMD, Cloudy sky with **light to moderate rainfall** may be expected from 22.06.2024 to 26.06.2024 in Chamarajanagara district. The day temperature is expected to be 28-31°C & night temperature is expected 16.4-18°C. The relative humidity in the morning hours is expected to be 79-86% & afternoon relative humidity is expected to be in the range of 49-68%. Wind speed expected to be 13-21 km/ hr.

**Recommendations to the farmers:**

Crop	Pest/Disease	Damage symptoms	Control measures
<b>Crops that can grown in the month of June</b>			
<b>Paddy:</b> BR-2655, Jaya, MSN-99			
<b>Aerobic Rice:</b> Daksha (KMP-175), Vanasiri (MAS-26), Sharada (MAS-946), Anaga (ARB-6), Rashi			
<b>Finger millet:</b> INDAF-8, MR-1, MR-6, L-5, KMR-301, GPU-45, KMAR-316			
<b>Kodo millet:</b> RBK-155, GPUK-3			
<b>Maize:</b> Hema, Nityashree, MAH-14-5			
<b>Shaktiman Maize:</b> CSH-5, CSH-9, CSV-4			
<b>Pop corn:</b> Amber			
<b>Little millet:</b> Co-2, OLM-203, GPUL-6			
<b>Barugu:</b> GPUP-8, GPUP-21, GPUP-28			
<b>Navane:</b> PS-4, SIA-326, GPUF-3			
<b>Barnyard millet:</b> DHBM-93-3			
<b>Bajra:</b> WCC-75, PHVB-910, ICTP-8203, SMV-221, MH-946			

**Groundnut:** GKVK-5, GKVK-27, KCG-6, GpBD-4, ICGV,-91114, JL-24, TMV-2  
**Sunflower:** KBSH-41, KBSH-42, KBSH-44, KBSH-53, KBSH-78, KBSH-85  
**Castor:** DSH-9, DSH-177, GSH-4, 48-1  
**Niger:** KBN-1  
**Red gram:** BRG-1, BRG-2, BRG-36, BRG-4, BRG-5, ICP-7035, TTB-7, Hyderabad-3C  
**Sugarcane:** Co-62175, Co-86032, CoVC-18061  
**Horticultural crops:** Tomato, Potato, Brinjal, Chilli, Cabbage, Radish, Knol Khol, Carrot, Onion, Mango, Banana, Lemon, Guava, Mosambi, Sapota, papaya, Pomegranate, Pineapple, Cashew nut, Garlic, Ginger, Tumeric  
**Fodder crops:**  
**Maize:** African tall  
**Sorghum:** MP Chaari, Pusa Chaari, JS-3, JS-20, COFS-29  
**Bajra:** Deena Bhandu-49A  
**Cowpea:** KBC-2, MFC-09-3

<b>Raising of seedlings in nursery beds (seedlings required per acre)</b>		
<b>Area</b>	:	<b>25 ft. length, 4 ft. wide and 4"height beds of 30 (300 sqm. /acre).</b>
<b>Seed</b>	:	<b>25 (kg./acre)</b>
<b>Selection of good quality seed</b>	:	<b>Add the seeds in water and common salt solution (1: 4 proportion) and mix well, half filled and Chaffey seeds will float, separate these seeds and wash good seeds in fresh water and dry under shade.</b>
<b>Land preparation</b>	:	<b>Plough the land properly, remove the weeds and make raised beds after levelling.</b>
<b>Nutrients required and sowing</b>	:	<b>For every 300 sqm. apply 750 kg FYM and 5.85 kg urea,8.4 kg super phosphate and 2.25kg muriate of potash or any equivalent inorganic fertilizers and mix well in the soil, sowing of treated seeds equally in all 30 nursery beds.</b>
<b>Water management</b>	:	<b>Thinly sprinkle the water two to three times per day.</b>
<b>Weed management</b>	:	<b>Within 24 hrs after sowing for every 300 sqm. nursery area, spray 75 ml Butachlor (50 E.C.) or 30 ml Pretelachlor + Saffner (30 E.C) solution in 25 ltr water or 300 Bensulphuron methyl (0.6% G.) + Pretelachlor (6.0% G.) or 7.5 g. Pyrozosulphuron ethyl (10% wp) mix with 5 - 10 kg sand and broadcast or Pretelachlor 120 g. or 3 or 4 days after of sowing, 24 ml. Anilophos (30 E.C.) in 25 ltr of water and subsequent weeds can be removed by hand weeding.</b>
<b>Plant protection</b>	:	<b>Spray 100 ml. Monocrotophos 36 S.L. after 10 - 12 days after sowing in 50 litres water against green leaf hopper.</b>
<b>Transplanting of seedlings</b>		<b>20 - 25 days after sowing</b>
<b>Seed treatment of Green gram, black gram, cowpea, Pigeon pea</b>		
<b>Treat the seeds with bio-fertilizers like Rhizobium and Phosphorus Solubilising Bacteria (PSB) @ 200 g. /acre, each.</b>		
<b>Maize</b>	<b>Fall army worm</b>	<b>5% damage- Application of 5% neem seed kernel extract (NSKE) or azadirachtin 1500ppm @ 5ml/litre (1 litre/acre)of water.</b>

		<b>10% damage- Whorl application of any one of the recommended insecticides for FAW, viz.,Chlorantraniliprole 18.5 SC (80 ml/acre) @ 0.4 ml/litre or Spinetoram 11.7 %SC (100ml/acre) @ 0.5 ml/litre or Emamectin benzoate 5% SG (80g/acre) @ 0.4g/litre.</b>
<b>Cowpea</b>	<b>Aphids</b>	<b>Several individuals suck the sap from tender shoots, flower buds, flowers, leaves and tender pods. Yellowing of leaves, shrivelled grains and sooty mold on leaves. To manage this pest spray Dimethoate – 30 EC. @ 1.7 ml./liter water spray. 200 lit. Spray mixture/acre.</b>
Tobacco	Tobacco leaf curl virus (Vector - Whitefly)	<ul style="list-style-type: none"> <li>• Remove and destroy the diseased tobacco seedlings before and after planting whenever they are seen in the field. Fill the gaps with healthy seedlings if it is not too late.</li> <li>• Alternate weed hosts for whitefly should be removed and destroyed.</li> <li>• In endemic areas <i>Sesbania</i> sp. may be grown as a barrier crop around tobacco fields.</li> <li>• Install 12 yellow sticky traps (castor oil coated) per hectare to monitor the whitefly population. If 100-200 whiteflies stick to the trap the following insecticide schedule has to be adopted.</li> <li>• 1<sup>st</sup> spray Chlorpyrifos 20% E.C. @ 25 ml. In 10 litres of water</li> <li>• 2<sup>nd</sup> spray Monocrotophos 36%, W.S.C. @ 15 ml in 10 litres of water</li> <li>• 3<sup>rd</sup> spray Acephate 75% S.P. @ 10 gm in 10 litres of water</li> <li>• 4<sup>th</sup> spray Demeton-s-methyl 25% E.C. @ 15 ml in 10 litres of water</li> </ul>
<b>Coconut</b>	<b>Rugose whitefly</b>	<ul style="list-style-type: none"> <li>✓ <b>Spraying starch solution (1%) to dislodge the heavy sooty mould deposition on the leaves of infested plants.</b></li> <li>✓ <b>Use of yellow sticky traps to trap the adult whiteflies</b></li> <li>✓ <b>In case of severe infestation, spray neem oil 0.5%</b></li> </ul>
<b>Black gram</b>	<b>Aphid</b>	<p><b>Adults and nymphs suck the sap from tender shoots, leaves; flower buds, flowers and tender pods. Yellowing of leaves, shriveled grain and development of sooty mold.</b></p> <p><b>Spray 1.7 ml. Dimethoate 30 EC. in lit. water, when infestation noticed. 250 lit. spray solution/acre.</b></p>
<b>Cowpea</b>	<b>Pod borer</b>	<p><b>Caterpillar bore the pods and feed on seeds. Infested pods with hole.</b></p> <p><b>On tender pods spray Chlorpyrifos - 20 EC. @ 2 ml./lit. water. 300 lit. spray solution/acre.</b></p>
<b>Chilli</b>	<b>Use of growth regulator</b>	<ul style="list-style-type: none"> <li>• <b>Generally flower drop will be noticed in chilli crop.</b></li> <li>• <b>To prevent flower drop spray NAA (Planofix) 50 ppm (50 mg. in 1 litre water) at time of flowering results in higher yield.</b></li> </ul>
<b>Tomato</b>	<b>Serpentine leaf miner</b>	<b>Larva mines the leaf epidermis, serpentine tunnels on leaves. Spray 0.3 ml. Imidachloprid 17.8 SL. or 2.0 ml. Triazophos 40 EC/lit. water.</b>
	<b>Fruit borer</b>	<ul style="list-style-type: none"> <li>• <b>Caterpillar bore the flower buds and fruits. Infested flower buds with hole and drops off, fruit with a hole, water enter through the hole leads to rotting.</b></li> </ul>

<b>Tomato</b>		<ul style="list-style-type: none"> <li>Trap crop: For every 25 rows of tomato grow one row of marigold cultivar African tall. The marigold seedlings about 35-40 days old. If borer problems exceeds 10 per cent spray 4 per cent. NSKE or 100 LE, Ha. NPV. If infestation in severe form spray 1.0 g. Methomil 40 SP. in a lit. water</li> </ul>
<b>Coconut</b>	<b>Rhinoceros beetle</b>	<b>Remove the beetle from infected part and fill 2 % Quinolphos or 5 % Melathion in sand @ 1:1 ratio.</b>
<b>Ginger</b>	<b>Rhizome treatment</b>	<b>Treat the rhizomes with 2 gm bleaching powder, 1 gm Metalaxyl MZ- 72, 1 gm Streptocycline mixed in 1 ltr of water. Treat the rhizomes 30 minutes and dry in shade. This will prevent rhizome rot problems.</b>
<b>Ginger</b>	<b>Rhizome rot</b>	<p><b>Symptoms: Yellowing of leaves, drying of leaf margin followed by burning of entire leaf, bottom portion of the shoot with water soaked soft, orange coloured rhizomes turn to brown colour, gradual rotting of such rhizomes.</b></p> <p><b>To manage this disease by drench the base of the plants with 2.0 g. Captan 50 WP or 2.0 g. Metalaxyl - MZ 72WP. In a lit.water.</b></p>

### Block level weather forecast (From 22-06-2024 to 26-06-2024)

#### Chamarajanagara

Parameter	22.06.2024	23.06.2024	24.06.2024	25.06.2024	26.06.2024
Rainfall (mm)	0	8.1	6.4	0.1	0.4
Max. temp (°C)	25.8	24	23.8	24	21.7
Min.Temp (°C)	17.7	17.7	18	17.9	17.8
Sky condition (Octas)	8	8	8	8	8
Relative humidity (%) 0830 hours	78	86	84	81	81
Relative humidity (%) 1730 hours	41	56	61	59	68
Wind Speed (kmph)	24	19	21	26	28
Wind Direction	248	248	248	248	248

#### Gundlupete

Parameter	22.06.2024	23.06.2024	24.06.2024	25.06.2024	26.06.2024
Rainfall (mm)	0	12.3	8.1	0.9	3.3
Max. temp (°C)	25.7	23.2	23.2	23	21.3
Min.Temp (°C)	17.6	17.7	17.8	17.7	17.7
Sky condition (Octas)	7	8	8	8	8
Relative humidity (%) 0830 hours	82	89	87	86	87
Relative humidity (%) 1730 hours	48	63	71	70	77

<b>Wind Speed (kmph)</b>	23	19	21	25	27
<b>Wind Direction</b>	248	248	248	248	248

<b>Kollegala</b>					
<b>Parameter</b>	<b>22.06.2024</b>	<b>23.06.2024</b>	<b>24.06.2024</b>	<b>25.06.2024</b>	<b>26.06.2024</b>
<b>Rainfall (mm)</b>	0	12.3	9.1	0	0
<b>Max. temp (°C)</b>	28.5	26.5	25.5	26.5	24.9
<b>Min.Temp (°C)</b>	20	19.7	19.8	20.2	20.1
<b>Sky condition (Octas)</b>	8	8	8	8	8
<b>Relative humidity (%) 0830 hours</b>	76	85	84	78	79
<b>Relative humidity (%) 1730 hours</b>	40	56	63	58	65
<b>Wind Speed (kmph)</b>	26	19	23	32	32
<b>Wind Direction</b>	248	248	248	248	248

<b>Yelandur</b>					
<b>Parameter</b>	<b>22.06.2024</b>	<b>23.06.2024</b>	<b>24.06.2024</b>	<b>25.06.2024</b>	<b>26.06.2024</b>
<b>Rainfall (mm)</b>	0	10.6	7.8	0.1	0.3
<b>Max. temp (°C)</b>	26.8	25	24.2	24.9	22.9
<b>Min.Temp (°C)</b>	18.6	18.6	18.7	18.8	18.7
<b>Sky condition (Octas)</b>	8	8	8	8	8
<b>Relative humidity (%) 0830 hours</b>	78	86	84	80	81
<b>Relative humidity (%) 1730 hours</b>	41	56	63	59	68
<b>Wind Speed (kmph)</b>	25	19	22	29	31
<b>Wind Direction</b>	248	248	248	248	248

- Download “**DAMINI**” app to get early warning on lightening and take precautions based on the alert given by the application.
- Kindly download”**MAUSAM**” APP for location specific forecast & warning &“**MEGHDOOT**” APP for Agromet advisory
- This information is available in the website: [mausam.imd.gov.in](http://mausam.imd.gov.in)

For any information farmers can contact **Dr.C.Ramachandra**, Senior Farm Superintendent/ **Dr. Sumanth Kumar.G.V**, Research Associate over phone No. 0821-259126/ 9535345814.

**AMFU of IMD,  
Naganahalli, Mysuru**