

**UNIVERSITY OF AGRICULTURAL SCIENCES, BENGALURU &  
INDIAN METEOROLOGICAL DEPARTMENT**



**GRAMIN KRISHI MAUSAM SEWA  
AMFU, OFRS, NAGANAHALLI,  
MYSURU - 570003**



**Date: 24-09-2024**

**AGRO-ADVISORY BULLETIN FOR MANDYA DISTRICT**

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

**Past Weather Data**

<b>Parameter</b>	<b>21.09.2024</b>	<b>22.09.2024</b>	<b>23.09.2024</b>	<b>24.09.2024</b>
<b>Rainfall (mm)</b>	0	0	0	0
<b>Max. Temp. (°C)</b>	32.6	31	33	33.6
<b>Min. Temp. (°C)</b>	19.9	18.7	20.2	-
<b>Sky condition (Octas)</b>	4	2	2	8
<b>Relative humidity (%) 0830 hours</b>	84	73	80	79
<b>Relative humidity (%) 1730 hours</b>	52	-	58	59
<b>Wind Speed (km/h)</b>	-	-	-	4
<b>Wind Direction</b>	-	-	-	230

**Weather forecast for the next five days (From 25-08-2024 to 29-09-2024)**

<b>Parameter</b>	<b>25.09.2024</b>	<b>26.09.2024</b>	<b>27.09.2024</b>	<b>28.09.2024</b>	<b>29.09.2024</b>
<b>Rainfall (mm)</b>	0	0	0	7	9
<b>Max. temp (°C)</b>	31.8	32.6	31.6	31.2	31
<b>Min.Temp (°C)</b>	17.3	17.2	17.3	17.6	18.6
<b>Sky condition (Octas)</b>	3	2	6	6	6
<b>Relative humidity (%) 0830 hours</b>	92	92	89	92	92
<b>Relative humidity (%) 1730 hours</b>	47	42	44	53	57
<b>Wind Speed (kmph)</b>	13	13	13	15	17
<b>Wind Direction</b>	248	283	257	249	249

**Forecast Summary**

As forecast received from IMD, cloudy sky with **light rainfall** may be expected from 25.09.2024 to 29.09.2024 in Mandya district. The day temperature is expected to be 31-32.6°C & night temperature is expected 17.2-18.6°C. The relative humidity in the morning hours is expected to be 89-92% & afternoon relative humidity is expected to be in the range of 42-57% per cent. Wind speed expected to be 13-17 km/ hr.

Recommendations to the farmers:			
Crop	Pest/Disease	Damage symptoms	Control measures
<b>Crops and varieties that can be grown in the month of August</b>			
<b>Finger millet :</b> Indaf-7, Indaf-9, KMR-301, GPU-45, KMR-316 <b>Paddy :</b> MSN-99 <b>Maize :</b> Hema, Nityashree, MAH-14-5 <b>Rabi Maize :</b> M-35-1, Nose (5-4-1), CSH-10 <b>Popcorn :</b> Amber <b>Sunflower:</b> KBSH-41, KBSH-42, KBSH-44, KBSH53, KBSH-78, KBSH-85 <b>Soybean:</b> MAUS-2 (Praja), Karune (Vegetable Soybean), KBS-23 <b>Niger:</b> KBN-1, No-71 <b>Cowpea :</b> TVK-944-02E, KBC-1, KBC-2, KBC-9, IT-98456-1, KM-5, KC-8 (K .BC-11) <b>Horse gram :</b> PHG-9, KBH-1 5209: 2.20-8371, 2.2.A.2-99463 (Vishal), VCF-0517 (Baahubali ), 222-18061 <b>Horticulture Crops:</b> Banana, Arecanut, Pineapple, Cauliflower, Onion <b>Fodder crops:</b> <b>Maize :</b> African Tall; <b>Maize:</b> MP Chari, Pusachari, JS-3, GS-20, COFS-29; <b>Bajra:</b> Dhina Bandhu- 49A; <b>Cowpea:</b> KBC-2			
<b>General recommendations for agricultural activities based on the given rainfall forecast:</b>			
<ul style="list-style-type: none"> <li>✓ Since there is light rainfall and rising temperatures, ensure timely irrigation for all crops, especially those in critical growth stages such as vegetative, flowering, and fruiting.</li> <li>✓ Drip irrigation or furrow irrigation can be employed to minimize water wastage and provide consistent moisture to the crops.</li> <li>✓ Apply organic mulches (like straw or dry leaves) around the base of crops to conserve soil moisture, reduce soil temperature, and prevent weed growth.</li> <li>✓ High temperatures can cause nutrient deficiencies. Monitor the crops and apply fertilizers based on soil testing to ensure healthy growth.</li> <li>✓ Foliar sprays of micronutrients can help alleviate nutrient stress caused by dry conditions.</li> <li>✓ Weed competition for water and nutrients should be minimized. Perform manual or chemical weeding based on the crop type.</li> <li>✓ With dry weather and high temperatures, monitor crops for pest infestations, such as sucking pests (aphids, whiteflies), which thrive in such conditions.</li> <li>✓ Use neem-based bio-pesticides or pheromone traps to control pests, and ensure proper field hygiene to minimize disease occurrence.</li> <li>✓ Use shading nets for heat-sensitive crops, especially vegetables, to reduce temperature stress and protect young plants from direct sunlight.</li> </ul>			
Crop	Stage	Weather-Based Advisory	
<b>Field Bean</b>	Harvesting	- Complete harvesting before rainfall on 28th and 29th Sept. to avoid quality loss.	
<b>Banana</b>	Bunch Development	- Support the plants with props to avoid lodging due to expected winds (up to 17 km/h). - Provide light irrigation until rainfall.	
<b>Paddy</b>	Vegetative Stage	- Ensure proper drainage during rainfall to avoid waterlogging. - Continue monitoring for pest and disease attacks.	
<b>Ragi</b>	Vegetative Stage	- Provide light irrigation until rainfall begins. - Maintain soil moisture and avoid water stress.	

<b>Red Gram</b>	Vegetative Stage	<ul style="list-style-type: none"> <li>- Apply organic mulch to conserve soil moisture.</li> <li>- Light irrigation before the rainfall can support growth.</li> </ul>
<b>Papaya</b>	Vegetative Stage	<ul style="list-style-type: none"> <li>- Support plants with stakes to avoid damage from strong winds.</li> <li>- Mulching around plants to conserve moisture is advisable.</li> </ul>
<b>Brinjal</b>	Fruiting Stage	<ul style="list-style-type: none"> <li>- Harvest mature fruits before 28th Sept. rainfall.</li> <li>- Ensure drainage to prevent fruit rot from excess moisture.</li> </ul>
<b>Chilli</b>	Flowering Stage	<ul style="list-style-type: none"> <li>- Avoid water stress; light irrigation is beneficial before expected rains.</li> <li>- Monitor for flower drop due to fluctuating moisture.</li> </ul>
<b>Cotton</b>	Boll Formation	<ul style="list-style-type: none"> <li>- Avoid waterlogging to prevent boll rot.</li> <li>- Support plants against possible winds on 28th and 29th Sept.</li> </ul>
<b>Coconut, Arecanut, Cocoa, Pepper</b>	Various Stages	<ul style="list-style-type: none"> <li>- Maintain mulch around trees for moisture conservation.</li> <li>- Inspect trees for pest infestations after rains.</li> </ul>
<b>Coffee</b>	Berry Development	<ul style="list-style-type: none"> <li>- Mulch and irrigation management until rainfall begins.</li> <li>- Monitor for berry borer after rainfall events.</li> </ul>
<b>Ginger</b>	Harvesting	<ul style="list-style-type: none"> <li>- If nearing maturity, harvest before the rainfall to avoid rhizome rot.</li> <li>- Dry the harvested crop in a covered area.</li> </ul>
<b>Sugarcane</b>	Vegetative Stage	<ul style="list-style-type: none"> <li>- Continue irrigation till rainfall occurs.</li> <li>- Ensure drainage during heavy rains to prevent root lodging.</li> </ul>
<b>Coconut black headed caterpillar</b>	Various stages	<ul style="list-style-type: none"> <li>• Remove and burn the severely affected fronds.</li> <li>• On community basis feed the Manocrotophos 36 SL. to the palm through root.</li> </ul> <p><b>Method:</b> A meter away from trunk, dig out and select brown coloured pencil thickness size root. Cut the root in a slanting position. To the polythene bag (size of 15 cm. length 4 cm. breadth) add 7.5 to 10 ml. Monocrotophos 36 SL. with equal quantity of water, introduce and immerse cut end of the root in insecticide mixture and tie the bag with thread.</p> <ul style="list-style-type: none"> <li>• The palm absorb the chemical within a period of 24 hours, if not after 48 hours select another root to feed the chemical.</li> <li>• A month after chemical treatment release larval parasites: gravid, Goniozus@ 10 - 12 /palm.</li> </ul> <p><b>Caution:</b> Not to harvest tender coconuts/matured coconuts for 30 days from date of chemical treatment.</p>
<b>Papaya mosaic ring spot virus</b>	Fruit development	<p>Nursery may be raised in 40 - 50 mesh nylon netting for a period of 60 days then plant.</p> <p>Around the garden 2 - 3 rows of African tall Maize should be grown on border crodiv. 30 - 40 days prior to papaya planting. Again after 2 months resowing of Maize by the side of previous Maize crodiv.</p> <p>Throughout the papaya cropping period maintain border crop of Maize.</p> <p>For control of sucking pests spray Dimethoate - 1.7 ml. /lit. water. Periodical spray is necessary.</p> <p>Note: June - July papaya planting can minimise the disease problem.</p> <p>Select disease free seedlings for planting.</p>

<b>Paddy Leaf folder</b>	Vegetative stage	Apply any one of the following insecticides per lit. water a) Quinalphos 25 EC. - 2.0 ml. b) Indoxacarb 14.5 SC. - 0.5ml. c) Flubendiamide 48 SC. - 0.08ml. d) Flubendiamide 20 WG. - 0.2 g. Drain out the water and spray the insecticide. 250 - 300 lit. spray mixture requires per acre.
<b>Red gram wilt</b>	Vegetative stage	5.0 g. Trichoderma viridae OR 3.0 g. Carbendazim + Mancozeb 75 WP.then sown. In wilt endemic areas before sowing enriched Trichoderma FYM incorporated to soil OR Sow wilt resistant red gram variety BRG 5 or Maruthi (ICP 8863).
<b>Paddy Yellow stem borer</b>	Vegetative stage	If infestation noticed, apply any one of the following insecticides per lit. water a) Monocrotophos 36 SL. - 1.5 ml. b) Chlorpyrifos 20 EC. - 2.0 ml. c) Flubendiamide 48 SC. - 0.08 ml. d) Flubendiamide 20 WG. - 0.2 g. Granular insecticide - kg./acre a) Fipronil 0.3 G - 10.0 b) Carbofuran 3 G - 8.0 N.B: Before application of granular insecticides, drain out the water and apply granules. Two days after application irrigate lightly.
<b>Coconut</b>	Rhinoceros beetle	Remove the adult beetle from crown of the palm by means of iron hook. Quinalphos 1.5 D. OR Malathion 5 D. mix with equal quantity of sand and plug the hole with mixture. Avoid FYM pits in and around coconut garden OR Mix 350 g.Quinalphos 1.5 D/ 3 m <sup>2</sup> of FYM.
<b>Paddy leaf and neck blast</b>	Transplanting to Vegetative	> Seed treatment: Treat the seeds @ 4 g. Carbendazim 50 WP. or Tricyclazole 75 WP. @ 0.6 g./kg. seed. Nursery spray > When seedlings are 10 -12 days old spray any one of the following fungicides to a lit. water. a) Carbendazim 50 WP. - 1.0 g. b) Tricyclazole 75 WP. - 0.6 g. c) Edifenphos 50 EC. - 1.0 ml. d) Kitazin 48 EC. - 1.0 ml. 20 - 25 days after transplanting if disease incidence above 5 per cent sprays any one fungicide mention above. If necessary spray at flowering stage. 200 - 300 lits. spray solution/acre.
<b>Coconut Eriophyid mites</b>	-	Addition to application of recommended NPK add 1 kg. Gypsum, 50 g. Boran, 5 kg. neem oil cake/palm. Spray 80 WP. Sulphur @ 4 g./lit. water on 2 - 6 months old tender nuts. Root feeding the mixture of 7.5 ml. Neemzol. OR

10 ml. Econeem with equal quantity of water.

### Poultry and Livestock

Category	Condition	Recommendation
Poultry	General	<ul style="list-style-type: none"> <li>• Use ventilation, exhaust fans, and sprinklers to cool the poultry house. Wet the roof or use a misting system to reduce heat.</li> <li>• Provide cool, clean water with electrolytes and vitamins (e.g., Vitamin C) to reduce heat stress.</li> <li>• Feed during early morning or late evening to avoid heat stress.</li> <li>• Litter Management: Keep litter dry to prevent ammonia build-up and respiratory issues.</li> </ul>
Livestock	General	<ul style="list-style-type: none"> <li>• Provide fresh, clean water and electrolyte solutions to avoid dehydration and heat stress.</li> <li>• Ensure shaded or ventilated shelters. Use fans or sprinklers in sheds to cool livestock.</li> <li>• Feed green fodder and silage. Avoid heat-generating feeds like excessive grains.</li> <li>• Monitor for signs of heat stress and deworm/vaccinate to prevent disease outbreaks.</li> </ul>

### Block level weather forecast (From 25-09-2024 to 29-09-2024)

#### Krishnarajpet

Parameter	25.09.2024	26.09.2024	27.09.2024	28.09.2024	29.09.2024
Rainfall (mm)	2.4	2	0.9	7.8	4.9
Max. temp (°C)	30.5	30.4	30.8	29.3	29.4
Min.Temp (°C)	19.3	18.8	19.7	20.1	19.3
Sky condition (Octas)	7	2	7	6	8
Relative humidity (%) 0830 hours	92	93	86	92	88
Relative humidity (%) 1730 hours	38	44	41	53	64
Wind Speed (kmph)	14	14	15	15	20
Wind Direction	248	249	252	257	249

#### Maddur

Parameter	25.09.2024	26.09.2024	27.09.2024	28.09.2024	29.09.2024
Rainfall (mm)	0.3	0.3	0.6	9.3	10.6
Max. temp (°C)	32.1	32.2	32	30.7	30.8
Min.Temp (°C)	20.9	20.2	21.5	21.1	20.7
Sky condition (Octas)	7	3	7	6	8
Relative humidity (%) 0830 hours	90	91	85	92	89
Relative humidity (%) 1730 hours	42	45	43	55	64

<b>Wind Speed (kmph)</b>	13	13	13	14	17
<b>Wind Direction</b>	248	248	252	252	248

<b>Malvalli</b>					
<b>Parameter</b>	<b>25.09.2024</b>	<b>26.09.2024</b>	<b>27.09.2024</b>	<b>28.09.2024</b>	<b>29.09.2024</b>
<b>Rainfall (mm)</b>	0.8	0.1	0.4	4.7	8.4
<b>Max. temp (°C)</b>	31	31.1	30.9	30	29.8
<b>Min.Temp (°C)</b>	20.1	19.6	21	20.3	20.1
<b>Sky condition (Octas)</b>	7	2	7	6	7
<b>Relative humidity (%) 0830 hours</b>	90	91	85	91	90
<b>Relative humidity (%) 1730 hours</b>	41	43	43	52	64
<b>Wind Speed (kmph)</b>	14	15	15	16	19
<b>Wind Direction</b>	248	248	249	249	248

<b>Mandya</b>					
<b>Parameter</b>	<b>25.09.2024</b>	<b>26.09.2024</b>	<b>27.09.2024</b>	<b>28.09.2024</b>	<b>29.09.2024</b>
<b>Rainfall (mm)</b>	0.4	3.3	0.4	6.7	8.4
<b>Max. temp (°C)</b>	31.4	31.5	30.7	29.9	30.3
<b>Min.Temp (°C)</b>	20.6	19.7	21.1	20.5	20.1
<b>Sky condition (Octas)</b>	7	3	8	6	7
<b>Relative humidity (%) 0830 hours</b>	89	91	84	91	89
<b>Relative humidity (%) 1730 hours</b>	40	46	45	54	64
<b>Wind Speed (kmph)</b>	15	14	14	15	18
<b>Wind Direction</b>	248	248	249	252	248

<b>Nagamangala</b>					
<b>Parameter</b>	<b>25.09.2024</b>	<b>26.09.2024</b>	<b>27.09.2024</b>	<b>28.09.2024</b>	<b>29.09.2024</b>
<b>Rainfall (mm)</b>	0.9	2.2	0.7	11.7	9.7
<b>Max. temp (°C)</b>	32	31.7	31.9	30	30.9
<b>Min.Temp (°C)</b>	19.9	19.1	20.1	20.5	19.7
<b>Sky condition (Octas)</b>	6	3	6	6	8
<b>Relative humidity (%) 0830 hours</b>	89	91	85	92	88
<b>Relative humidity (%) 1730 hours</b>	40	46	44	58	64
<b>Wind Speed (kmph)</b>	14	14	15	15	19
<b>Wind Direction</b>	249	283	283	257	249

**Pandavapura**

<b>Parameter</b>	<b>25.09.2024</b>	<b>26.09.2024</b>	<b>27.09.2024</b>	<b>28.09.2024</b>	<b>29.09.2024</b>
<b>Rainfall (mm)</b>	1	3.1	0.5	7	6.8
<b>Max. temp (°C)</b>	30.6	30.3	30.4	29.1	29.2
<b>Min.Temp (°C)</b>	20	19.2	20.5	20	19.7
<b>Sky condition (Octas)</b>	7	3	7	6	8
<b>Relative humidity (%) 0830 hours</b>	90	92	85	92	89
<b>Relative humidity (%) 1730 hours</b>	39	47	44	53	64
<b>Wind Speed (kmph)</b>	15	15	15	15	19
<b>Wind Direction</b>	248	248	249	249	248

**Shrirangapattana**

<b>Parameter</b>	<b>25.09.2024</b>	<b>26.09.2024</b>	<b>27.09.2024</b>	<b>28.09.2024</b>	<b>29.09.2024</b>
<b>Rainfall (mm)</b>	0.6	3.1	0.3	6.5	7.4
<b>Max. temp (°C)</b>	29.9	29.8	29.4	28.5	28.5
<b>Min.Temp (°C)</b>	19.7	18.8	20.2	19.5	19.3
<b>Sky condition (Octas)</b>	7	3	7	6	8
<b>Relative humidity (%) 0830 hours</b>	90	92	85	93	90
<b>Relative humidity (%) 1730 hours</b>	39	47	44	54	65
<b>Wind Speed (kmph)</b>	15	15	16	16	19
<b>Wind Direction</b>	248	248	249	249	248


- Download “**DAMINI**” app to get early warning on lightening and take precautions based on the alert given by the application.
- Kindly download “**MAUSAM**” APP for location specific forecast & warning & “**MEGHDOOT**” APP for Agromet advisory
- This information is available in the website: [mausam.imd.gov.in](http://mausam.imd.gov.in)

For any information farmers can contact **Dr. C. Ramachandra**, Senior Farm Superintendent/ **Dr. Sumanth Kumar.G.V**, Technical officer over phone No. 0821-259126/ 9535345814.

**AMFU of IMD,**  
**Naganahalli, Mysuru**

वास्तविक वर्षा तथा विस्तारित अवधि पूर्वानुमान  
 Realized Rainfall and Extended Range Forecast  
 (वर्षा और तापमान)  
 (Rainfall and Temperature)


Realized Rainfall  
 (5<sup>th</sup> to 18<sup>th</sup> September, 2024)





## Extended Range Forecast System

### Rainfall forecast maps for the next 2 weeks (IC- 18<sup>th</sup>September, 2024) (20<sup>th</sup>September to 03<sup>rd</sup> October, 2024)




- **Week1 (20.09.2024 to 26.09.2024):** Rainfall is likely to be normal in parts of Northeast India and Central India. Below normal rainfall is likely over East India, Northwest India, Himachal Pradesh, Uttarakhand, Uttar Pradesh, Konkan & Goa, Karnataka and Kerala.
- **Week 2 (27.09.2024 to 03.10.2024):** Rainfall is likely to be above normal over most parts of the country. Rainfall is likely to be normal in Northeast India and Tamil Nadu.

**Maximum and Minimum temperature anomaly ( °C) forecast  
for the next 2 weeks (IC- 18<sup>th</sup>September, 2024)  
(20<sup>th</sup>September to 03<sup>rd</sup> October, 2024)**

**MME forecast Tmax anomaly (Deg C)**

(Week1: 20Sep–26Sep)

(Week2: 27Sep–03Oct)




**Maximum Temperature (Tmax)**

- **Week 1 (20.09.2024 to 26.09.2024):** Maximum temperature is likely to be above normal over most parts of the country.
- **Week 2 (27.09.2024 to 03.10.2024):** Maximum temperature is likely to be above normal over Jammu & Kashmir, Himachal Pradesh, Uttarakhand, East Uttar Pradesh, Bihar and Northeast India.

**MME forecast Tmin anomaly (Deg C)**

(Week1: 20Sep–26Sep)

(Week2: 27Sep–03Oct)



**Minimum Temperature (Tmin)**

- **Week 1 (20.09.2024 to 26.09.2024) and Week 2 (27.09.2024 to 03.10.2024):** Tmin is likely to be above normal in most parts of Northwest India, Central India and Karnataka. Tmin is likely to be below normal Eastern coastal states and Kerala.

