

**UNIVERSITY OF AGRICULTURAL SCIENCES, BENGALURU &  
INDIAN METEOROLOGICAL DEPARTMENT**



**GRAMIN KRISHI MAUSAM SEWA  
AMFU, OFRS, NAGANAHALLI,  
MYSURU - 570003**



Date:26-11-2024

**AGRO-ADVISORY BULLETIN FOR MYSURU DISTRICT**

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

**Past Weather Data**

<b>Parameter</b>	<b>22.11.2024</b>	<b>23.11.2024</b>	<b>24.11.2024</b>	<b>25.11.2024</b>	<b>26.11.2024</b>
<b>Rainfall (mm)</b>	0	0	0	-	0
<b>Max. Temp. (°C)</b>	29.6	31.2	29.9	-	28.6
<b>Min. Temp. (°C)</b>	17.9	18.2	18	-	18.5
<b>Sky condition (Octas)</b>	3	3	0	-	7
<b>Relative humidity (%) 0830 hours</b>	74	76	71	-	75
<b>Relative humidity (%) 1730 hours</b>	131	62	24	-	-
<b>Wind Speed (km/h)</b>	4	192	0	-	6
<b>Wind Direction</b>	50	250	0	-	50

**Weather forecast for the next five days (From 27-11-2024 to 01-12-2024)**

<b>Parameter</b>	<b>27.11.2024</b>	<b>28.11.2024</b>	<b>29.11.2024</b>	<b>30.11.2024</b>	<b>01.12.2024</b>
<b>Rainfall (mm)</b>	0	0	0	0	0
<b>Max. temp (°C)</b>	23.9	24.5	25.8	26.2	25.7
<b>Min.Temp (°C)</b>	17.9	17.7	17.5	18.2	18.4
<b>Sky condition (Octas)</b>	8	7	7	8	8
<b>Relative humidity (%) 0830 hours</b>	90	81	80	88	90
<b>Relative humidity (%) 1730 hours</b>	60	50	51	52	56
<b>Wind Speed (kmph)</b>	5.9	6.6	5.6	5.8	1.1
<b>Wind Direction</b>	38	29	15	4	18

**Forecast Summary**

As forecast received from IMD, partially cloudy sky with **no rainfall** may be expected from 27.11.2024 to 01.12.2024 in Mysuru district. The day temperature is expected to be 23.5-26.2°C & night temperature is expected 17.5-18.4°C. The relative humidity in the morning hours is expected to be 80-90% & afternoon relative humidity is expected to be in the range of 51-60%. Wind speed expected to be 1.1-6.6 km/hr.

**SMS Advisory**

Protect crops and livestock from cold; irrigate late morning and provide warm shelters for animals.

**Recommendations to the farmers:-**

Crop	Pest/Disease	Damage symptoms	Control measures
<b>General Advisory:</b>			
<ul style="list-style-type: none"> <li>✓ As no rainfall is expected, schedule light but consistent irrigation for crops like cabbage, cauliflower, tomato, and beans.</li> <li>✓ Avoid over-irrigation, particularly for crops in fruit and pod development stages, to prevent diseases like fruit rot and root rot.</li> <li>✓ Monitor crops regularly for pests such as aphids, pod borers, and fruit borers.</li> <li>✓ Use eco-friendly pest control methods such as neem oil or pheromone traps.</li> <li>✓ Mulch around crops to conserve soil moisture and suppress weed growth.</li> <li>✓ Apply balanced fertilizers, especially potash and nitrogen, to support growth in critical stages like fruiting and pod formation.</li> <li>✓ Prune dead or diseased parts to promote healthy growth.</li> <li>✓ Protect sensitive crops like banana and coffee from strong winds by supporting them with stakes.</li> <li>✓ For turmeric and ginger ready for harvest, ensure proper drying of rhizomes to prevent fungal growth.</li> <li>✓ Mulch and maintain basin formation around coconut, arecanut, and black pepper to conserve soil moisture.</li> <li>✓ Provide clean, dry shelters and adequate ventilation to livestock.</li> <li>✓ Increase feeding of high-energy fodder to maintain body warmth in cooler temperatures.</li> <li>✓ Maintain optimal room temperature (24-26°C) and humidity (65-75%) for silkworm rearing.</li> <li>✓ Feed silkworms fresh and healthy mulberry leaves to ensure uniform growth.</li> <li>✓ Ensure poultry houses are well-ventilated and dry.</li> <li>✓ Provide clean drinking water and balanced feed to maintain productivity.</li> <li>✓ Look for early signs of fungal infections due to high humidity during the morning.</li> <li>✓ Use appropriate fungicides or organic solutions like garlic extracts for management.</li> <li>✓ Avoid field operations during peak midday hours to prevent heat exhaustion.</li> <li>✓ Ensure proper storage of harvested produce to maintain quality.</li> </ul>			

**Weather based advisory**

Crop	Stage	Advisory
<b>Cabbage and cauliflower</b>	<b>Head formation stage</b>	Maintain adequate soil moisture through light irrigation. Watch for pests like aphids.
<b>Bean</b>	<b>Pod formation stage</b>	Ensure consistent soil moisture. Handpick pests like pod borers if observed.
<b>Tomato</b>	<b>Fruit development stage</b>	Stake plants to prevent fruit contact with soil. Avoid overwatering to prevent diseases.
<b>Red gram</b>	<b>Pod initiation stage</b>	Monitor for pod borers. Apply a light dose of fertilizers for healthy pod development.
<b>Paddy</b>	<b>Hard dough stage</b>	Avoid waterlogging. Monitor for pests and prepare for harvesting soon.
<b>Chilli</b>	<b>Fruit development</b>	Remove damaged fruits and monitor for fruit rot or viral

	<b>stage</b>	infections.
<b>Field bean</b>	<b>Pod development</b>	Irrigate moderately. Monitor for pod pests like aphids.
<b>Banana</b>	<b>Fruit development stage</b>	Support the plants with props to prevent lodging. Apply potash-rich fertilizers.
<b>Chilli</b>	<b>Vegetative stage</b>	Perform timely weeding. Apply nitrogen fertilizers to promote healthy vegetative growth.
<b>Horticultural crops</b>	<b>Various stages</b>	Monitor for pests/diseases. Adjust irrigation based on stage-specific needs.
<b>Plantation crops</b>	<b>Various stages</b>	Mulch around the base to conserve moisture. Prune older leaves to promote airflow.
<b>Livestock</b>	<b>Shelter and Feeding</b>	Provide clean, dry shelters. Maintain hydration and balanced feed to support health.
<b>Sericulture</b>	<b>Rearing stage</b>	Maintain optimal temperature and humidity in rearing rooms. Feed silkworms with fresh mulberry leaves.
<b>Poultry</b>	Shelter and Feeding	Ensure adequate ventilation in coops. Provide clean water and balanced feed.

### Recommendation to farmers

#### Crop specific advisory:

<b>Crop</b>	<b>Stage</b>	<b>Advisory</b>
<b>Cabbage diamond back moth</b>	Head stage	<ul style="list-style-type: none"> <li>• Spray DDVP 76 EC. @0.5 ml./lit water in nursery.</li> <li>• 15 days before transplanting around the main field and every 25 rows of cabbage one row of mustard sowing, 15 to 20 days after cabbage planting another row of mustard sowing. Mustard as trap crop. Spray on mustard with 0.5 ml. DDVP in a lit. water.</li> <li>• During head formation, spray 5 per cent NSKE .</li> <li>• Birdpurchases may be provided to attract predatory birds.</li> </ul>
<b>Tomato whiteflies</b>	Fruiting stage	Spray 1.0ml.Oxydemeton methyl 25 EC in a lit. water.
<b>Bean Pod borer</b>	Pod formation stage	Spray 2.0 ml. Malathion 50 EC./ lit. water .
<b>Tomato Early and late blight of tomato</b>	Fruiting stage	<p>For late blight of tomato 15 days prior to transplanting Trichoderma and Pseudomonas enriched compost may be incorporated to the soil.  For early blight control spray 2.0 g. Mancozeb 75 WP  OR  2.0 g. Maneb  OR  2.0 g. Metalaxyl- MZ 72WP.  OR  2.0 g. Dimethomorph + polyram/lit. water.  For control of late blight spray 2.0 g. Metalaxyl - MZ 72WP.  OR</p>

		2.0 g. Fosetyl al 80 WP OR 2.0 g. Dimethomorph + polyram in a lit. water, 5 weeks after transplanting. Repeat the spray 7th, 9th and 11th weeks after transplanting. 200- 250 lit. spray solution required/acre/spray.
<b>Rice earhead bug</b>	Hard dough stage	> During milky stage of the crop; spray Malathion 50 EC. at 2.0 ml./lit. of water .  <b>OR</b> > Dust 8 - 10 kg. Malathion 5 D./acre during morning hours.
<b>Rice Brown plant hoppers</b>	Hard dough stage	Spray any one of the following insecticides per lit. water 1) Imidacloprid 17.8 SL.- 0.5 ml. 2) Thiamethoxam 25 WG.- 0.7 g. 3) Monocrotophos 36 SL.- 1.5ml 4) Chlorpyriphos 20 EC.- 2.0 ml. 5) Buprofezin 25 EC.- 1.4ml. > Spray solution should reach the base of the plant. > Around 400 to 450 lit. spray solution required/acre. Granular insecticide kg./ac 1) Carbofuran 3 G- 8.0 2) Phorate 10 G- 5.0 3) Quinalphos 5 G - 12.0 N.B: Drain out the water and apply granules. Two days after application light irrigation may be provided.
<b>Red gram Sterility mosaic</b>	Pod initiation stage	Pull out the infested plants and destroy. 20 - 25, 40 - 45 days after sowing spray 2.5 ml. Dicofol 18.5 EC./lit. water. ICP 7035 sterility mosaic resistant red gram variety.
<b>Banana Leaf spot (Cigatoka)</b>	Fruit development	In endemic areas grow resistant banana variety - Sakkare bale. At the time of planting the rhizomes may treated with any one of the Fungicides /lit. water a)Propiconazole 25 EC.- 1.0 ml. b)Theiophenate methyl 70 Wdiv.- 1.0 g. c)Carbendazim 50 Wdiv.- 1.0 g. d)Metham Sodium (Vapom) - 1.0 g. In Mashy area provide drainage.
<b>Field bean pod borer</b>	Pod development	Dust 10 kg. Fenvalrate 0.4 D. OR Malathion 5 D. per acre during morning hours.
<b>Paddy Leaf folder</b>	Panicle emergence stage	Apply any one of the following insecticides per lit. water a) Quinalphos 25 EC. - 2.0 ml. b) Indoxacarb 14.5 SC. - 0.5ml. c) Flubendiamide 48 SC. - 0.08ml. d) Flubendiamide 20 WG. - 0.2 g. Drain out the water and spray the insecticide. 250 - 300 lit. spray mixture requires per acre.
<b>Paddy Bacterial leaf blight</b>	Panicle emergence stage	25 and 50 DAT add 0.5 g. Streptocycline and 2.5 g. Copper oxychloride 50 WP for a lit. Water and spray. 200 to 250 lit. Spray mixture requires/acre/time.

<b>Ginger Rhizome rot</b>	Harvesting stage	2.0 g. Metalaxyl - MZ 72Wdiv. in a lit. water. Before store of seed material soak them in 3.0 g. Mancozeb 75 Wdiv. in a lit. water for 30 min then dry in shade and store.
---------------------------	------------------	--

**Block level weather forecast (From 27-11-2024 to 01-12-2024)**

<b>H.D. Kote</b>					
<b>Parameter</b>	<b>27.11.2024</b>	<b>28.11.2024</b>	<b>29.11.2024</b>	<b>30.11.2024</b>	<b>01.12.2024</b>
<b>Rainfall (mm)</b>	0	0	0	0	0
<b>Max. temp (°C)</b>	25.5	24.5	25.6	26.4	26.9
<b>Min.Temp (°C)</b>	18.6	18.2	16.9	17.1	18.5
<b>Sky condition (Octas)</b>	8	7	7	8	7
<b>Relative humidity (%) 0830 hours</b>	92.4	84.2	79.7	79.1	92.4
<b>Relative humidity (%) 1730 hours</b>	57.4	57.6	48.1	54.7	53.7
<b>Wind Speed (kmph)</b>	6.9	7	4.6	4.4	1.9
<b>Wind Direction</b>	51.3	34.5	18.4	350.6	291.8

<b>Hunsuru</b>					
<b>Parameter</b>	<b>27.11.2024</b>	<b>28.11.2024</b>	<b>29.11.2024</b>	<b>30.11.2024</b>	<b>01.12.2024</b>
<b>Rainfall (mm)</b>	0	0	0	0	0
<b>Max. temp (°C)</b>	25.2	24.2	25.1	25.8	25.9
<b>Min.Temp (°C)</b>	18.1	18.2	16.7	16.8	18.4
<b>Sky condition (Octas)</b>	8	8	7	8	8
<b>Relative humidity (%) 0830 hours</b>	91.8	82	79.3	78.7	87.1
<b>Relative humidity (%) 1730 hours</b>	57.3	57.4	48.3	55.4	53.9
<b>Wind Speed (kmph)</b>	7.9	9.4	8.8	8.3	4.3
<b>Wind Direction</b>	43.1	22.6	11.7	355.1	335.6

<b>K.R. Nagara</b>					
<b>Parameter</b>	<b>27.11.2024</b>	<b>28.11.2024</b>	<b>29.11.2024</b>	<b>30.11.2024</b>	<b>01.12.2024</b>
<b>Rainfall (mm)</b>	0	0	0	0	0
<b>Max. temp (°C)</b>	25.2	24.2	25.1	25.6	25.7
<b>Min.Temp (°C)</b>	18.2	18.2	16.9	17	18.2
<b>Sky condition (Octas)</b>	8	7	7	8	8

<b>Relative humidity (%) 0830 hours</b>	92.1	81.4	77.6	77.5	84.2
<b>Relative humidity (%) 1730 hours</b>	57.1	56	46.9	53.8	53.3
<b>Wind Speed (kmph)</b>	8.4	9	8.8	8.4	5.8
<b>Wind Direction</b>	43.2	23.5	11.7	352.6	338.2
<b>Mysuru</b>					
<b>Parameter</b>	<b>27.11.2024</b>	<b>28.11.2024</b>	<b>29.11.2024</b>	<b>30.11.2024</b>	<b>01.12.2024</b>
<b>Rainfall (mm)</b>	0	0	0	0	0
<b>Max. temp (°C)</b>	25.5	24.6	25.2	26.4	26.1
<b>Min.Temp (°C)</b>	18.9	18.8	17.5	17.5	18.8
<b>Sky condition (Octas)</b>	8	7	7	8	8
<b>Relative humidity (%) 0830 hours</b>	92.6	79.8	75.8	80.2	90.7
<b>Relative humidity (%) 1730 hours</b>	56	54.5	46.8	51.5	54.5
<b>Wind Speed (kmph)</b>	8.4	7.9	6.5	6.6	5.2
<b>Wind Direction</b>	50.2	24.2	90	347.5	326.3

<b>Nanjanagudu</b>					
<b>Parameter</b>	<b>27.11.2024</b>	<b>28.11.2024</b>	<b>29.11.2024</b>	<b>30.11.2024</b>	<b>01.12.2024</b>
<b>Rainfall (mm)</b>	0	0	0	0	0
<b>Max. temp (°C)</b>	25.7	24.2	25	26.4	26.2
<b>Min.Temp (°C)</b>	19	18.9	17.5	17.8	19
<b>Sky condition (Octas)</b>	8	7	7	8	8
<b>Relative humidity (%) 0830 hours</b>	91.6	81.3	76.9	81.5	92.9
<b>Relative humidity (%) 1730 hours</b>	55.1	57.8	49.9	52.1	57.1
<b>Wind Speed (kmph)</b>	6.5	4.3	1.8	3.3	5.4
<b>Wind Direction</b>	56.3	24.4	323.2	310.6	277.6

<b>Piriapatna</b>					
<b>Parameter</b>	<b>27.11.2024</b>	<b>28.11.2024</b>	<b>29.11.2024</b>	<b>30.11.2024</b>	<b>01.12.2024</b>
<b>Rainfall (mm)</b>	0	0	0	0	0
<b>Max. temp (°C)</b>	25	23.8	24.8	25.6	25.9
<b>Min.Temp (°C)</b>	17.5	17.6	16.2	16.4	17.8
<b>Sky condition (Octas)</b>	8	7	7	8	8
<b>Relative humidity (%) 0830 hours</b>	92.3	85.6	82.6	82	90.6

<b>Relative humidity (%) 1730 hours</b>	57.6	58.8	49	56.8	55.7
<b>Wind Speed (kmph)</b>	7.6	9.1	8.2	7.6	3.7
<b>Wind Direction</b>	45	33.7	23.2	5.4	331

### T. Narasipura

<b>Parameter</b>	<b>27.11.2024</b>	<b>28.11.2024</b>	<b>29.11.2024</b>	<b>30.11.2024</b>	<b>01.12.2024</b>
<b>Rainfall (mm)</b>	0	0	0	0	0
<b>Max. temp (°C)</b>	25.8	24.7	25.2	26.6	26.1
<b>Min.Temp (°C)</b>	19	18.9	17.7	17.8	19.1
<b>Sky condition (Octas)</b>	8	7	7	8	8
<b>Relative humidity (%) 0830 hours</b>	90.6	78.6	76.8	79	89.9
<b>Relative humidity (%) 1730 hours</b>	56.7	56	49.1	52.3	58
<b>Wind Speed (kmph)</b>	6.9	6.3	4	3.6	4.7
<b>Wind Direction</b>	47.1	13.2	333.5	323.2	292.6


- Download “**DAMINI**” app to get early warning on lightening and take precautions based on the alert given by the application.
- Kindly download “**MAUSAM**” APP for location specific forecast & warning & “**MEGHDOOT**” APP for Agromet advisory
- This information is available in the website: [mausam.imd.gov.in](http://mausam.imd.gov.in)

For any information farmers can contact **Dr.C.Ramachandra**, Senior Farm Superintendent/ **Dr. Sumanth Kumar.G.V**, Technical officer over phone No.0821-259126/ 9535345814.

**AMFU of IMD,  
Naganahalli, Mysuru**

वास्तविक वर्षा तथा विस्तारित अवधि पूर्वानुमान  
**Realized Rainfall and Extended Range Forecast**  
 (वर्षा और तापमान)  
 (Rainfall and Temperature)


**Realized Rainfall**  
 (07<sup>th</sup> to 20<sup>th</sup> November, 2024)






## Extended Range Forecast System

**Rainfall forecast maps for the next 2 weeks (IC- 20<sup>th</sup> November, 2024)  
(22<sup>nd</sup> November to 05<sup>th</sup> December, 2024)**



- **Week1 (22.11.2024 to 28.11.2024):** Rainfall is likely to be normal over South India.
- **Week 2 (29.11.2024 to 05.12.2024):** Rainfall is likely over South India, Central India, East India and Nagaland, Manipur, Mizoram & Tripura (NMMT).

**Maximum and Minimum temperature anomaly (°C) forecast  
for the next 2 weeks (IC- 20<sup>th</sup> November, 2024)  
(22<sup>nd</sup> November to 05<sup>th</sup> December, 2024)**




### Maximum Temperature (Tmax)

- **Week 1 (22.11.2024 to 28.11.2024):** Maximum temperature is likely to be above normal over Jammu & Kashmir, Punjab, Himachal Pradesh, West Rajasthan and Chhattisgarh. It

is likely to be below normal over Central India, West India and South India.

- **Week 2 (29.11.2024 to 05.12.2024):** Maximum temperature is likely to be above normal over Jammu & Kashmir, Himachal Pradesh, Punjab, West Rajasthan and Arunachal Pradesh. It is likely to be below normal over Central India, West India, South India and some parts of East India.



#### Minimum Temperature (Tmin)

- **Week 1 (22.11.2024 to 28.11.2024):** Minimum temperature is likely to be below normal over most parts of the country.
- **Week 2 (29.11.2024 to 05.12.2024):** Minimum temperature is likely to be below normal over many parts of Northwest India, Central India and some parts of East India. It is likely to be above normal over Jammu & Kashmir, Northeast India, many parts of West India and South India.