# UNIVERSITY OF AGRICULTURAL SCIENCES, BENGALURU & INDIAN METEOROLOGICAL DEPARTMENT



## GRAMIN KRISHI MAUSAM SEWA AMFU, OFRS, NAGANAHALLI, MYSURU - 570003



Date:14-03-2025

#### AGRO-ADVISORY BULLETIN FOR KODAGU DISTRICT

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

Past Weather Data							
Parameter	11.03.2025	12.03.2025	13.03.2025	14.03.2025			
Rainfall (mm)	0	0	0	0			
Max. Temp. (°C)	33.3	35.7	34.8	33.9			
Min. Temp. (°C)	19.4	20.4	18.2	19.2			
Sky condition (Octas)	-	-	ı	-			
Relative humidity (%) 0830 hours	100	99	90	72			
Relative humidity (%) 1730 hours	78	68	65	23			
Wind Speed (km/h)	-	-	-	-			
Wind Direction	-	-	-	-			

## Weather forecast for the next five days (From 15-03-2024 to 19-03-2025)

Parameter	15.03.2025	16.03.2025	17.03.2025	18.03.2025	19.03.2025
Rainfall (mm)	0	0	0	0	2
Max. Temp. (°C)	35	35	35	35	35
Min. Temp. (°C)	18	18	18	18	18
Sky condition (Octas)	3	3	2	3	3
Relative humidity (%) 0830 hours	85	79	77	82	79
Relative humidity (%) 1730 hours	29	29	30	37	30
Wind Speed (kmph)	10	10	8	8	8
Wind Direction	334	90	45	140	225

#### **Forecast Summary**

As forecast received from IMD, cloudy sky with very light rainfall may be expected from 15.03.2025 to 19.03.2025 in Kodagu district. The day temperature is expected to be 35°C & night temperature is expected 18°C. The relative humidity in the morning hours is expected to be 77-85% & afternoon relative humidity is expected to be in the range of 29-37%. Wind speed expected to be 8-10 km/hr.

#### **SMS Advisory**

A forecasted temperature for the next five days is 35°C. Farmers should irrigate crops adequately and use mulching to conserve soil moisture. Provide shade and sufficient drinking water for livestock to prevent heat stress. Very light rainfall aids in minimizing heat-related damage for crops.

Recommendations to the farmers:-						
Crop	Pest/Disease	Damage symptoms	Control measures			
General Advisory	7:					

- Very light rainfall prevents soil moisture loss, despite providing irrigation at proper intervals is essential to avoid water stress.
- Mulching with organic materials (leaves, straw, dried grass) is advised to retain soil moisture and reduce evaporation.
- Drip and sprinkler irrigation are recommended for efficient water management.
- **Pest and Disease Monitoring**: The dry weather can increase **thrips**, **mites**, **aphids**, **and borers**; regular scouting and preventive control measures should be taken.
- **Shade Management** in plantation crops like coffee and cardamom is necessary to protect plants from direct sunlight and water stress.

Crop	Stage	Weather-Based Advisory
Paddy	Nursery	Very light rainfall: despite of very light rainfall ensure adequate
	preparation/transp	irrigation for nursery. Maintain proper moisture for seedling growth.
7.5.	lanting	
Maize	Tasseling stage	Very light rainfall; aids in preventing water stress. Mulching is
		recommended to conserve soil moisture. Monitor for fall
D :	<b>37</b>	armyworm.
Ragi	Vegetative stage	Very light rainfall reduces moisture loss.
Tomato	Fruit development	Very light rainfall: Install mulch to maintain soil moisture. Protect
D	TI : /C :.:	from fruit borers and powdery mildew.
Brinjal	Flowering/fruitin	Very light rainfall: avoids flower drop. Monitor for sucking pests
G1 1111	g stage	like whiteflies and mites.
Chilli	Flowering stage	Very light rainfall: despite irrigate regularly to avoid flower drop.
G 11	TT 1C	Monitor for thrips and mites due to dry conditions.
Cabbage	Head formation	<b>Very light rainfall:</b> Regularly check for diamondback moth infestation.
Cauliflower	Head formation	Very light rainfall: keeps the soil moist for better head
		development. Monitor for aphids and caterpillars.
Coffee	Berry	Very light rainfall: provides protective irrigation. Mulching and
	development	shade tree management are recommended. Avoid excessive
	(post-harvest	pruning.
	maintenance)	
Arecanut	Fruiting/maturity	Very light rainfall: supports nut filling. Irrigate at 10-15 days
		interval. Protect from mites and spindle bug infestations.
Coconut	Nut development	Very light rainfall: apply irrigation to prevent nut drop. Use
		organic mulches to retain soil moisture. Keep palms free from
		rugose whitefly.
Pepper	Fruiting	Very light rainfall: maintain soil moisture through light irrigation.
		Provide shade and mulch. Check for pollu beetle infestations.
Cardamom	Maintenance/post	Very light rainfall: helps in moisture retention through mulching.
	-harvest phase	Maintain shade trees.
Banana	Bunch	Very light rainfall: avoids fruit shrinkage. Provide propping
	development	support to avoid lodging due to dry winds.
Sapota	Fruiting/harvest	Very light rainfall: is essential in maintaining fruit quality. Timely
		harvesting is advised to prevent fruit cracking.
Guava	Fruiting	Very light rainfall; aids to maintain fruit size and quality.
		Mulching can be beneficial.

Livestock, Poultry, and Sericulture Advisory (Very light Rainfall & High Temperature					
Sector	Weather-Based Advisory				
Livestock	Ensure proper shade and ventilation in animal sheds. Provide ample clean drinking				
	water. Avoid grazing during peak heat hours. Provide mineral supplements to prevent				
	heat stress.				
Poultry	High temperatures may lead to heat stress, affecting egg production and bird health.				
	Maintain proper ventilation in poultry sheds. Provide cool drinking water with				
	electrolytes. Reduce feed quantity in the daytime and provide more during cooler				
	hours.				
Sericulture	High temperatures can stress silkworms. Maintain humidity by sprinkling water in				
	rearing rooms. Provide proper aeration and shade to protect mulberry plants from heat				
	stress.				

Moisture Conservation Practices and Summer Ploughing Advisory					
Practice	Weather-Based Advisory				
Mulching	Apply dry leaves, paddy straw, or organic waste around plants to reduce				
	evaporation losses and soil temperature.				
Summer Ploughing	Since rainfall is absent, conduct deep summer ploughing to expose soil-borne				
	pests and improve aeration. It also helps in better moisture retention for the				
	next season.				
Irrigation	Follow drip irrigation or sprinkler irrigation to conserve water. Irrigate during				
Management	early morning or evening hours to minimize evaporation losses.				
<b>Shading Measures</b>	For young plants and nurseries, use shade nets or temporary structures to				
	reduce direct heat impact.				

Block level weather forecast (From 15-03-2025 to 19-03-2025)							
Madikeri							
Parameter 15.03.2025 16.03.2025 17.03.2025 18.03.2025 19.03.2025							
Rainfall (mm)	4.9	7.3	0	0	0		
Max. temp (°C)	32.6	30.2	33.5	33.7	34.1		
Min.Temp (°C)	19.1	19.8	19.7	19.7	19.4		
Sky condition (Octas)	6	8	5	4	2		
Relative humidity (%) 0830 hours	90.3	92.2	94.2	95	92.1		
Relative humidity (%) 1730 hours	33.4	43.3	31.9	30.6	24.9		
Wind Speed (kmph)	7.4	3.5	2.3	1.8	1.6		
Wind Direction	67.2	66	18.4	323.2	333.5		

Somvarpet						
Parameter	15.03.2025	16.03.2025	17.03.2025	18.03.2025	19.03.2025	
Rainfall (mm)	3	3.8	0	0	0	
Max. temp (°C)	32.4	30.9	33.9	34.4	34.4	
Min.Temp (°C)	18.4	19.6	19.8	19.5	19.4	
Sky condition (Octas)	6	7	6	4	4	
Relative humidity (%) 0830 hours	90.4	90.8	93	95	93	
Relative humidity (%) 1730 hours	30.7	37.9	26.1	27.2	22.6	
Wind Speed (kmph)	7.9	3.6	1.5	1.8	0.7	
Wind Direction	87.4	110	346	281.3	110	

Virajpet						
Parameter	15.03.2025	16.03.2025	17.03.2025	18.03.2025	19.03.2025	
Rainfall (mm)	2.8	7.3	0	0	0	
Max. temp (°C)	34.6	32.1	35.1	34.9	35.2	
Min.Temp (°C)	20.5	20.8	20.9	21	20.6	
Sky condition (Octas)	6	8	6	3	3	
Relative humidity (%) 0830 hours	87.9	94.2	97.3	98	94.6	
Relative humidity (%) 1730 hours	36.6	44.2	39.5	41.6	32.5	
Wind Speed (kmph)	8	3.9	2.4	1.4	1.5	
Wind Direction	54.1	56.3	26.5	90	14	

- Download "DAMINI" app to get early warning on lightening and take precautions based on the alert given by the application.
- ➤ Kindly download"MAUSAM"APP for location specific forecast & warning &"MEGHDOOT" APP for Agromet advisory
- > This information is available in the website: mausam.imd.gov.in


For any information farmers can contact **Dr. C. Ramachandra**, Senior Farm Superintendent/ **Dr. SumanthKumar.G.V**, Technical officer over phone No. 0821-259126/ 9535345814

AMFU of IMD,

Naganahalli, Mysuru


# वास्तविकवर्षातथाविस्तारितअवधिपूर्वानुमान Realized Rainfall and Extended Range Forecast (वर्षाऔरतापमान)

(Rainfall and Temperature)




#### **Extended Range Forecast System**

Rainfall forecast maps for the next 2 weeks (IC- 12<sup>th</sup>March,2025) (14<sup>th</sup>to 27<sup>th</sup>March, 2025)



- Week1(14.03.2025 to 20.03.2025):Rainfall is likely to be above normalover North East India. Rainfall activity is also likely over Jammu & Kashmir, Himachal Pradesh, Uttarakhand, Kerala, coastal Karnataka and coastal regions of Tamil Nadu.
- Week 2 (21.03.2025 to 27.03.2025):Rainfall is likely to be above normalover Kerala, Tamil Nadu and Karnataka.Rainfall activity is also likely over Jammu & Kashmir,Himachal Pradesh, Uttarakhand, North East India, Maharashtra, Telanganaand some parts of Chhattisgarh and Odisha.


Maximum and Minimum temperature anomaly (°C) forecast for the next 2 weeks (IC- 12<sup>th</sup>March,2025) (14<sup>th</sup>to 27<sup>th</sup>March, 2025)



**Maximum Temperature (Tmax)** 

• Week 1 (14.03.2025 to 20.03.2025): Maximum temperature is likely to be above normal

- over most parts of North West India, East India, Konkan-Goa, Karnataka and States along East coast and below normal over Central India.
- Week 2 (21.03.2025 to 27.03.2025): Maximum temperature is likely to be above normal over most parts of North West India and many parts of East India. However, it is likely to be below normal over Central India, West India and South India.



## **Minimum Temperature (Tmin)**

- Week 1 (14.03.2025 to 20.03.2025): Minimum temperature is likely to be below normal over Central India, Jammu & Kashmir and Odisha. However, it is likely to be above normal over many parts of North West India, North East India, Gujarat, Madhya Maharashtra and Karnataka.
- Week 2 (21.03.2025 to 27.03.2025): Minimum temperature is likely to be below normal over most parts of the country. However, it is likely to be above normal over East Uttar Pradesh, Gujarat, Karnataka, Madhya Maharashtraand some parts of North East India.